




Draft Genomic Sequences of *Streptomyces misionensis* ACT66 and *Streptomyces albidoflavus* ACT77, Bacteria with Potential Application for Phytopathogen Biocontrol

 Victor Satler Pylro,^a Armando Cavalcante Franco Dias,^b Fernando Dini Andreote,^c Alessandro de Mello Varani,^d Cristiane Cipolla Fasanella Andreote,^b Iron Amoreli de Figueiredo Ribeiro,^e Isabella Takahashi Kitano,^e Eduardo Roberto de Almeida Bernardo^e

^aDepartment of Biology, Federal University of Lavras (UFLA), Lavras, Minas Gerais, Brazil

^bAndrios Assessoria, Piracicaba, São Paulo, Brazil

^cSoil Microbiology Laboratory, Luiz de Queiroz College of Agriculture, University of São Paulo (ESALQ/USP), Piracicaba, São Paulo, Brazil

^dDepartamento de Tecnologia, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista (Unesp), Jaboticabal, São Paulo, Brazil

^eAgrivalle Brasil Indústria e Comércio de Produtos Agrícolas, Ltda, Salto, São Paulo, Brazil

ABSTRACT Here, we report the draft genomic sequences and annotation of *Streptomyces misionensis* ACT66 and *Streptomyces albidoflavus* ACT77, which are two bacteria with potential application for phytopathogen biocontrol.

Phytopathogen biocontrol using bacteria is a promising and eco-friendly strategy for replacing or reducing the application of agrochemicals (1). The knowledge of genomic features may help us to understand the mechanisms involved in the interaction between biological control agents and the target phytopathogen, which is crucial for enhancing the use of these organisms in agriculture. *Streptomyces misionensis* strain ACT66 and *Streptomyces albidoflavus* strain ACT77 are Gram-positive bacteria commonly found in soil, as are the other *Actinobacteria*. Preliminary analyses have indicated that these isolates show *in vitro* fungicidal activity, suggesting their potential application for phytopathogen biocontrol *in situ* (data not shown). To gain insight into the use of these bacteria for biological control of phytopathogens, we performed whole-genome sequencing (WGS). Several *Streptomyces* spp. have been used to promote plant growth and for phytopathogen biocontrol, for example, against *Magnaporthe oryzae* in rice (2), *Fusarium* spp. (3), and some wood decay fungi (WDF) (4). *S. misionensis* strain ACT66 and *S. albidoflavus* strain ACT77 were isolated from soil under agriculture management in Brazil, by the dilution plating technique onto 5% tryptone soy agar (TSA; Bacto BD, USA) supplemented with 50 mg ml⁻¹ of benomyl. The plates were incubated at 28°C, and the isolates were kept as pure cultures. To perform the WGS, the isolates were cultivated in nutrient agar for 48 h, at 28°C, and the genomic DNA was extracted using the Wizard genomic DNA purification kit (Promega) following the manufacturer's instructions.

Paired-end sequencing libraries (2 × 250 bp) were constructed using the Nextera XT kit (Illumina, San Diego, CA) following the manufacturer's instructions and sequenced using the Illumina MiSeq platform (Illumina). After quality filtering using Trimmomatic version 0.33 (5) (parameters [paired-end reads] included trailing, 10; leading, 10; slidingwindow, 4:10), a total of 1,185,642 paired-end reads were obtained for *S. misionensis* strain ACT66, and 540,058 paired-end reads were obtained for *S. albidoflavus* strain ACT77, consisting of a genome coverage of ~70× and ~36×, respectively. All reads were reference-based assembled with SPAdes version 3.12 (6), using *S. misionensis* strain DSM 40306 (GenBank accession number [NZ_FNTD000000000](https://doi.org/10.1128/MRA.00949-19)) for *S. misionensis* ACT66 and *S. albidoflavus* strain NRRL B-1271 ([NZ_JOII000000000](https://doi.org/10.1128/MRA.00949-19)) for *S. albidoflavus*

Citation Pylro VS, Dias ACF, Andreote FD, Varani ADM, Andreote CCF, Ribeiro IADF, Kitano IT, Bernardo ERDA. 2019. Draft genomic sequences of *Streptomyces misionensis* ACT66 and *Streptomyces albidoflavus* ACT77, bacteria with potential application for phytopathogen biocontrol. *Microbiol Resour Announc* 8:e00949-19. <https://doi.org/10.1128/MRA.00949-19>.

Editor Frank J. Stewart, Georgia Institute of Technology

Copyright © 2019 Pylro et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Victor Satler Pylro, victor.pylro@ufla.br, or Eduardo Roberto de Almeida Bernardo, eduardo.bernardo@agrivalle.com.br.

Received 8 August 2019

Accepted 13 August 2019

Published 5 September 2019

strain ACT77 as references. The obtained contigs were further processed with the SIS software (7) to generate a set of contig scaffolds representing the draft genomes. The REAPR pipeline (8) was used to improve the assembly accuracy. Default parameters were used for all software unless otherwise noted.

This assembly procedure resulted in 210 scaffolds for *S. misionensis* strain ACT66 and 95 scaffolds for *S. albidoflavus* strain ACT77. Genome completeness and contamination were estimated using CheckM (9) in the lineage-specific mode. The estimated genome size for *S. misionensis* strain ACT66 is 8,312,220 bp, with a G+C content of 72.3% and an N_{50} value of 70,083 bp. The estimated genome size for *S. albidoflavus* strain ACT77 is 7,446,125 bp, with a G+C content of 73.2% and an N_{50} value of 1,152,190 bp. The genome completeness estimated by CheckM was 98.28% and 99.89%, and the contamination was 0.45% and 1.17% for *S. misionensis* strain ACT66 and *S. albidoflavus* strain ACT77, respectively, and they were classified as nearly complete with low contamination. We applied the method proposed by Parks and colleagues (10), which uses the software GTDBk and the Genome Taxonomy Database (GTDB; <http://gtdb.ecogenomic.org>) for assigning taxonomy to each assembled genome using the default parameters. Based on this software, our two isolates were classified as *S. misionensis* and *S. albidoflavus*.

Genome annotation was performed with PATRIC version 3.5.23 (11). It identified 7,936 coding DNA sequences (CDS) and 86 predicted noncoding RNAs (68 tRNAs and 18 rRNAs, encompassing 6 rRNA operons) for *S. misionensis* strain ACT66 and 6,819 CDS and 89 predicted noncoding RNAs (68 tRNAs and 21 rRNAs, encompassing 7 rRNA operons) for *S. albidoflavus* strain ACT77.

Data availability. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession numbers [VOGW00000000](https://www.ncbi.nlm.nih.gov/nuclseq/VOGW00000000) and [VOGX00000000](https://www.ncbi.nlm.nih.gov/nuclseq/VOGX00000000) for *Streptomyces misionensis* strain ACT66 and *Streptomyces albidoflavus* strain ACT77, respectively. The versions described in this paper are the first versions. Raw reads are available under the BioProject accession number [PRJNA557451](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA557451).

ACKNOWLEDGMENTS

This work was supported by the Brazilian Microbiome Project (<http://www.brmicrobiome.org>). V.S.P. received funds from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) grant number 404651/2018-6.

REFERENCES

- Canova SP, Petta T, Reyes LF, Zucchi TD, Moraes LA, Melo IS. 2010. Characterization of lipopeptides from *Paenibacillus* sp. (IIRAC30) suppressing *Rhizoctonia solani*. *World J Microbiol Biotechnol* 26:2241–2247. <https://doi.org/10.1007/s11274-010-0412-9>.
- Law JW, Ser HL, Khan TM, Chuah LH, Pusparajah P, Chan KG, Goh BH, Lee LH. 2017. The potential of Streptomyces as biocontrol agents against the rice blast fungus, *Magnaporthe oryzae* (*Pyricularia oryzae*). *Front Microbiol* 8:3. <https://doi.org/10.3389/fmicb.2017.00003>.
- Bubici G. 2018. *Streptomyces* spp. as biological control agents against *Fusarium* species. *CAB Rev Perspect Agric Vet Sci Nutr Nat Resour* 13:50.
- Jung SJ, Kim NK, Lee DH, Hong SI, Lee JK. 2018. Screening and evaluation of *Streptomyces* species as a potential biocontrol agent against a wood decay fungus, *Gloeophyllum trabeum*. *Mycobiology* 46:138–146. <https://doi.org/10.1080/12298093.2018.1468056>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Dias Z, Dias U, Setubal JC. 2012. SIS: a program to generate draft genome sequence scaffolds for prokaryotes. *BMC Bioinformatics* 13:96. <https://doi.org/10.1186/1471-2105-13-96>.
- Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD. 2013. REAPR: a universal tool for genome assembly evaluation. *Genome Biol* 14:R47. <https://doi.org/10.1186/gb-2013-14-5-r47>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil PA, Hugenholtz P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life. *Nat Biotechnol* 36:996. <https://doi.org/10.1038/nbt.4229>.
- Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, Gillespie JJ, Gough R, Hix D, Kenyon R, Machi D, Mao C, Nordberg EK, Olson R, Overbeek R, Pusch GD, Shukla M, Schulman J, Stevens RL, Sullivan DE, Vonstein V, Warren A, Will R, Wilson MJC, Seung Yoo H, Zhang C, Zhang Y, Sobral BW. 2014. PATRIC, the bacterial bioinformatics database and analysis resource. *Nucleic Acids Res* 42:D581–D591. <https://doi.org/10.1093/nar/gkt1099>.