



VALÉRIA FONSECA MOSCARDINI

**SUBLETHAL EFFECTS OF INSECTICIDAL
SUNFLOWER SEED TREATMENTS ON THREE
SPECIES OF APHID NATURAL ENEMIES**

LAVRAS – MG

2015

VALÉRIA FONSECA MOSCARDINI

**SUBLETHAL EFFECTS OF INSECTICIDAL SUNFLOWER SEED
TREATMENTS ON THREE SPECIES OF APHID NATURAL ENEMIES**

Tese apresentada à Universidade Federal de Lavras, como parte das exigências do Programa de Pós-Graduação em Entomologia, área de concentração em Entomologia, para a obtenção do título de Doutora.

Orientador

Dr. Geraldo Andrade Carvalho

LAVRAS – MG

2015

**Ficha catalográfica elaborada pelo Sistema de Geração de Ficha Catalográfica da Biblioteca
Universitária da UFLA, com dados informados pelo(a) próprio(a) autor(a).**

Moscardini, Valéria Fonseca.

Sublethal effects of insecticidal sunflower seed treatments on
three species of aphid natural enemies / Valéria Fonseca

Moscardini. – Lavras : UFLA, 2015.

72 p.

Tese(doutorado)–Universidade Federal de Lavras, 2015.

Orientador(a): Geraldo Andrade de Carvalho.

Bibliografia.

1. Parasitoid. 2. Extrafloral nectar. 3. Systemic insecticides. I.
Universidade Federal de Lavras. II. Título.

VALÉRIA FONSECA MOSCARDINI

**SUBLETHAL EFFECTS OF INSECTICIDAL SUNFLOWER SEED
TREATMENTS ON THREE SPECIES OF APHID NATURAL ENEMIES**

**(EFEITOS SUBLETAIS DE TRATAMENTO DE SEMENTES DE
GIRASSOL COM INSETICIDAS SOBRE TRÊS ESPÉCIES DE
INIMIGOS NATURAIS DE AFÍDEOS)**

Tese apresentada à Universidade Federal de Lavras, como parte das exigências do Programa de Pós-Graduação em Entomologia, área de concentração em Entomologia, para a obtenção do título de Doutora.

APROVADA em 12 de fevereiro de 2015.

Dr. J.P. Michaud - Kansas State University - EUA

Dr. Maurício Sérgio Zacarias – Embrapa - Café

Dr. Renê Luis de Oliveira Rigitano - UFLA

Dr. Ronald Zanetti Bonetti Filho - UFLA

Dr. Geraldo Andrade Carvalho

Orientador

LAVRAS – MG

2015

Aos meus pais Iara e Renato pelo amor, exemplo e incentivo,

DEDICO

AGRADECIMENTOS

A Deus pelo milagre da vida e por estar sempre presente em meu caminho.

À Universidade Federal de Lavras e ao Departamento de Entomologia pela oportunidade concedida para a realização do Doutorado.

À Fundação de Amparo à Pesquisa do estado de Minas Gerais (FAPEMIG), pela concessão da bolsa de estudos e pesquisa.

À Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) pela concessão de bolsa de Doutorado sanduíche.

Ao professor Dr. Geraldo Andrade Carvalho, pela orientação, amizade, confiança e pelos ensinamentos que foram de grande importância para realização deste trabalho e para meu crescimento profissional. Meus agradecimentos também a sua esposa Ana Paula, pelo agradável convívio e amizade.

Ao professor Dr. J.P. Michaud, pela amizade, paciência, orientação e ensinamentos de grande importância durante o Doutorado sanduíche.

À Kansas State University, Agricultural Research Center - Hays, pela oportunidade para realização do Doutorado sanduíche.

Ao meu esposo Pablo Gontijo, pela amizade, companheirismo, paciência, amor e pela inestimável parceria nos trabalhos realizados.

Aos amigos do laboratório de Seletividade Rafaella, Rodrigo, Jader, Thaís, Brenda, Mariana, Andrea, DeJane, Wellington, Lucas, Dyrson e Frontino pela amizade.

Às amigas Lilian, Fernanda, Elisangela, Ana Luiza e Érika, pela amizade, companheirismo, longas conversas e momentos de alegria.

Aos professores do Departamento de Entomologia da UFLA, pelos ensinamentos e harmoniosa convivência durante o doutorado.

Aos funcionários do Departamento de Entomologia da UFLA Lisiane, Érica, Elaine, Nazaré, e em especial Léa, D. Irene e Julinho, pela valiosa colaboração e amizade.

Aos amigos do laboratório da Kansas State University, Clint e Mohamed, pela valiosa ajuda nos trabalhos realizados, amizade e harmoniosa convivência.

Às minhas amigas bibliotecárias americanas, Paula Elder e Jean Curtis-Neitz, por disponibilizarem tempo para ensinar inglês, pela amizade, momentos de alegria e diversão.

Aos professores Dr. Renê Luis de Oliveira Rigitano, Dr. Maurício Sérgio Zacarias, Dr. Ronald Zanetti Bonetti Filho por participarem da banca examinadora e pelas sugestões.

Aos meus pais, Iara e Renato, pelo amor incondicional e incentivo, meus irmãos Vinícius e Viviani, pela amizade e companheirismo, aos meus avós Hebe, Alaíde (*in memoriam*) e Djalma, pela torcida e incentivo, aos meus sobrinhos pelos momentos de alegria, descontração e carinho, aos meus cunhados Thiago e Rafaella, pela amizade e apoio e a todos meus familiares que contribuíram durante toda essa jornada.

Aos meus sogros Selma e João, pelo carinho, amizade e incentivo, ao meu cunhado Douglas e concunhada Larissa pelos momentos de alegria e amizade e a todos das famílias Costa e Gontijo por me receberem com muito carinho.

A todos, que direta ou indiretamente, contribuíram para o êxito deste trabalho, os meus sinceros agradecimentos.

GENERAL ABSTRACT

The extrafloral nectar (EFN) of sunflower, *Helianthus annuus* L., is an important resource for many insects and represents a potential route of exposure to systemic insecticides applied as seed treatments to cultivated varieties. This study examined whether chlorantraniliprole or thiamethoxam might negatively impact *Lysiphlebus testaceipes*, *Coleomegilla maculata* and *Hippodamia convergens* when they consumed the EFN of sunflowers grown from treated seed. Consumption of EFN by *L. testaceipes* caused no lethal effects, but reduced the numbers of *Schizaphis graminum* Rondani attacked and parasitized in Petri dish arenas. Whereas control females self-superparasitized every fourth host, those exposed to chlorantraniliprole did not. Thiamethoxam greatly reduced the proportion of female offspring, suggesting an impact on the egg fertilization process. Exposure of *C. maculata* larvae to chlorantraniliprole and thiamethoxam delayed adult emergence by prolonging the pupal period. When adults were exposed, thiamethoxam reduced the preoviposition period compared to chlorantraniliprole, whereas the latter treatment caused females to produce fewer clutches. Larvae of *C. maculata* did not appear to obtain sufficient hydration from the sunflower stems and their subsequent fecundity and fertility were compromised. Exposure of *H. convergens* larvae to thiamethoxam skewed the sex ratio in favor of females; both materials reduced the egg viability of resulting adult females and increased the period required for eclosion. Exposure of *H. convergens* adults to chlorantraniliprole reduced egg eclosion times compared to thiamethoxam and exposure to both insecticides reduced pupation times in progeny.

Keywords: Parasitoid. *Coleomegilla maculata*. *Hippodamia convergens*. Extrafloral nectar. Systemic insecticides.

RESUMO GERAL

Néctar extrafloral (NEF) de girassol, *Helianthus annuus* L. é uma importante fonte de alimento para muitos insetos e apresenta-se como potencial via de exposição a inseticidas sistêmicos utilizados no tratamento de sementes para diversas culturas. Este estudo teve como objetivo avaliar se clorantraniliprole ou tiametoxam podem afetar negativamente *Lysiphlebus testaceipes*, *Coleomegilla maculata* e *Hippodamia convergens* quando estes insetos alimentaram-se de NEF de girassol provenientes de sementes tratadas. O consumo de NEF por adultos de *L. testaceipes* não causou efeito letal, mas reduziu o número de *Schizaphis graminum* Rondani atacados e parasitados quando expostos em placa de Petri aos adultos do parasitoide. Fêmeas do tratamento controle auto-superparasitaram um a cada quatro hospedeiros encontrados quando comparadas com fêmeas expostas a clorantraniliprole. Tiametoxam reduziu consideravelmente a proporção de progênie fêmeas. A exposição de larvas de *C. maculata* a clorantraniliprole e tiametoxam prolongou o período de pupa. Quando adultos foram expostos, tiametoxam reduziu o período de pré-oviposição em comparação com clorantraniliprole. Fêmeas expostas a clorantraniliprole tiveram número reduzido de ovos. Larvas de *C. maculata* possivelmente não conseguiram se alimentar de forma adequada das hastes de girassol e adultos provenientes destas larvas expostas apresentaram baixa fecundidade e fertilidade. Adultos oriundos de larvas de *H. convergens* expostas a tiametoxam tiveram a razão sexual alterada; ambos inseticidas reduziram a viabilidade dos ovos da geração subsequente e aumentaram o período de eclosão dos ovos. Exposição de adultos de *H. convergens* a clorantraniliprole reduziu o tempo de eclosão das larvas quando comparado com tiametoxam, e a exposição a ambos inseticidas reduziu o tempo de pupa na progênie.

Palavras-chave: Parasitoide. *Coleomegilla maculata*. *Hippodamia convergens*. Néctar extrafloral. Inseticidas sistêmicos.

SUMMARY

FIRST PART	10
GENERAL INTRODUCTION	10
REFERENCES	13
SECOND PART - ARTICLES	18
ARTICLE 1 Sublethal effects of chlorantraniliprole and thiamethoxam seed treatments when <i>Lysiphlebus testaceipes</i> feed on sunflower extrafloral nectar	18
ABSTRACT	19
INTRODUCTION	20
MATERIALS AND METHODS	22
RESULTS	26
DISCUSSION	27
REFERENCES	32
ARTICLE 2 Sublethal effects of insecticide seed treatments on two nearctic lady beetles (Coleoptera: Coccinellidae)	43
ABSTRACT	44
INTRODUCTION	45
MATERIALS AND METHODS	48
RESULTS	53
DISCUSSION	54
REFERENCES	60

FIRST PART

GENERAL INTRODUCTION

Over the past several years, systemic insecticides have been used as seed treatments in sunflower (*Helianthus annuus* L.), cotton (*Gossypium* spp.), oilseed rape (*Brassica napus* L.), corn (*Zea mays* L.), soybean (*Glycine max* L.) and other cereals and crops (BRADSHAW; RICE; HILL, 2008; STRAUSBAUGH; EUJAYL; FOOTE, 2010) to control pests that cause damage early in the crop (HODGSON; KEMIS; GEISINGER, 2012; NUYTTENS et al., 2013). Systemic insecticides used in the treatment of seeds are characterized by low lipophilicity (CLOYD; BETHKE, 2011) which facilitates their translocation in plant tissues.

Chlorantraniliprole and thiamethoxam represent two different systemic groups, diamides and neonicotinoids, respectively. The anthranilic diamide class of insecticides activates ryanodine receptors, thus stimulating uncontrolled calcium ion release from muscle cells, causing insect paralysis (CORDOVA et al., 2006; LAHM et al., 2007). In contrast, neonicotinoids target nicotinic acetylcholine receptors in the insect central nervous system, causing receptor blockage, paralysis and death (TOMIZAWA; CASIDA, 2005). Furthermore, the conversion of thiamethoxam into highly toxic metabolites such as clothianidin (CASIDA, 2011), a process which can occur within both plant (CLOYD; BETHKE, 2011) and insect (NAUEN et al., 2003) tissues, may increase their toxicity to insects and their natural enemies. Nauen et al. (2003) demonstrated that clothianidin is one of the primary metabolites of thiamethoxam in true leaves of cotton plants treated via soil drench. However, its greater lipophilicity and lower solubility in water make clothianidin more likely than thiamethoxam to contaminate nectar.

Seed treatment may reduce the total amount of insecticide applied by confining it to the seed and thus confer environmental advantages compared with broadcast or soil or applications (TAYLOR; ECKENRODE; STRAUB, 2001) including reduced exposure of beneficial insects (CLOYD; BETHKE, 2011; MIZELL; SCONYERS, 1992). Despite these advantages, research has revealed lethal and sublethal effects of seed treatments on non-target organisms such as *Orius insidiosus* (Say) (Hemiptera: Anthocoridae) (GONTIJO et al., 2014a), *Chrysoperla carnea* (Neuroptera: Chrysopidae) (GONTIJO et al., 2014b) and *Anagrus nilaparvatae* (Pang and Wang) (Hymenoptera: Mymaridae) (LIU et al., 2010). Negative effects are due contamination of pollen and nectar, both floral and extrafloral (EFN), which are often important food sources for natural enemies (CHOATE; LUNDGREN, 2013; LUNDGREN, 2009; WÄCKERS; ROMEIS; VAN RIJN, 2007).

Non-target effects of systemic insecticides on beneficial species are a substantial risk because the majority of crops, including commercial sunflower, are now planted with seed treatments. On the High Plains of the USA, sunflower plants are an important source of EFN for beneficial insects during the hot, dry summers (ROYER; WALGENBACH, 1991). Sunflower extrafloral nectaries are extremely small and highly abundant along petioles and leaf veins; they secrete EFN continuously from the time the first true leaves expand until the plants senesce. The production of EFN by plants is associated with attraction of herbivore natural enemies and thus fosters mutualistic protection for the plant (MARAZZI; BRONSTEIN; KOPTUR, 2013). Intercropping plants with EFN-producing plants has even been proposed to supplement food for natural enemies early in the growing season before pests become abundant (JAMONT; CREPELLIERE; JALOUX, 2013). The generalist aphid parasitoid, *Lysiphlebus testaceipes* (Cresson) (Hymenoptera: Braconidae) and two species of predatory lady beetle, *Coleomegilla maculata* DeGeer and *Hippodamia convergens*

Guérin-Méneville (Coleoptera: Coccinellidae), are among the beneficial species attracted to sunflower EFNs on the High Plains of the USA.

Lysiphlebus testaceipes is a solitary endoparasitoid that attacks many aphid species (PIKE et al., 2000). It is an important natural enemy of the greenbug, *Schizaphis graminum* (Rondani) (Hemiptera: Aphididae) and contributes to biological control of this aphid in both sorghum (JONES, 2001) and wheat (GILES et al., 2003). Without access to nectar or some other sugar source, many parasitoids show a dramatic reduction in their ability to parasitize their hosts (LEWIS et al., 1998; STAPEL et al., 1997). Lady beetles such as *C. maculata* and *H. convergens* also consume pollen and nectar, both floral and EFN (PEMBERTON; VANDENBERG, 1993; SMITH; KRISCHIK, 1999). These species are two of the most abundant coccinellids in the central USA, and *H. convergens* is of particular importance in providing biological control of cereal aphids in wheat, sorghum and other grains (MICHAUD, 2013; NECHOLS; HARVEY, 1998; RICE; WILDE, 1988).

Considering the economic importance of sunflowers in grain and oil production and the widespread use of insecticidal seed treatments, the objective of the present study was to assess whether chlorantraniliprole or thiamethoxam could have any negative or sublethal impacts on *L. testaceipes*, *C. maculata* or *H. convergens* when various life stages fed on the EFN of sunflower seedlings grown from treated seed.

REFERENCES

BRADSHAW, J. D.; RICE, M. E.; HILL, J. H. Evaluation of management strategies for bean leaf beetles (Coleoptera: Chrysomelidae) and bean pod mottle virus (Comoviridae) in soybean. **Journal of Economic Entomology**, Lanham, v. 101, n. 4, p. 1211-1227, aug. 2008.

CASIDA, J. E. Neonicotinoid metabolism: compounds, substituents, pathways, enzymes, organisms, and relevance. **Journal of Agricultural and Food Chemistry**, Washington, v. 59, n. 7, p. 2923-2931, apr. 2011.

CHOATE, B. A.; LUNDGREN, J. G. Why eat extrafloral nectar? Understanding food selection by *Coleomegilla maculata* (Coleoptera: Coccinellidae). **Biocontrol**, Gz Dordrecht, v. 58, n. 58, p. 359-367, jun. 2013.

CLOYD, R. A.; BETHKE, J. A. Impact of neonicotinoid insecticides on natural enemies in greenhouse and interiorscape environments. **Pest Management Science**, Hoboken, v. 67, n. 1, p. 3-9, aug. 2011.

CORDOVA D. et al. Anthranilic diamides: a new class of insecticides with a novel mode of action, ryanodine receptor activation. **Pesticide Biochemistry and Physiology**, San Diego, v. 84, n. 3, p. 196-214, mar. 2006.

GILES, K. L. et al. Development of a sampling plan in winter wheat that estimates cereal aphid parasitism levels and predicts population suppression. **Journal of Economic Entomology**, Lanham, v. 96, n. 3, p. 975-982, mar. 2003.

GONTIJO, P. C. et al. Non-target effects of two sunflower seed treatments on *Orius insidiosus* (Hemiptera: Anthocoridae). **Pest Management Science**, Hoboken, 2014a. doi:10.1002/ps.3798a

GONTIJO, P. C. et al. Non-target effects of chlorantraniliprole and thiamethoxam on *Chrysoperla carnea* when employed as sunflower seed treatments. **Journal of Pest Science**, Heidelberg, v. 87, n. 4, p. 711-719, dec. 2014b.

HODGSON, E. W.; KEMIS, M.; GEISINGER, B. Assessment of Iowa soybean growers for insect pest management practices. **Journal of Extension**, Ames, v. 50, n. 4, RIB6, aug. 2012.

JAMONT, M.; CREPELLIERE, S.; JALOUX, B. Effect of extrafloral nectar provisioning on the performance of the adult parasitoid *Diaeretiella rapae*. **Biological Control**, San Diego, v. 65, n. 2, p. 271-277, may. 2013.

JONES, D. B. **Natural enemy thresholds for greenbug, *Schizaphis graminum* Rondani, on winter wheat**. 2001. M.S. thesis - Oklahoma State University, Stillwater, USA, 2001.

LAHM, G. P. et al. Rynaxypyr: a new insecticidal anthranilic diamide that acts as a potent and selective ryanodine receptor activator. **Bioorganic and Medicinal Chemistry Letters**, New York, v. 17, n. 22, p. 6274-6279, sep. 2007.

LEWIS, W. J. et al. Understanding how parasitoids balance food and host needs: importance to biological control. **Biological Control**, San Diego, v. 11, n. 2, p. 175-183, feb. 1998.

LIU, F. et al. Effects of imidacloprid on the orientation behavior and parasitizing capacity of *Anagrus nilaparvatae*, an egg parasitoid of *Nilaparvata lugens*. **BioControl**, Gz Dordrecht, v. 55, n. 4, p. 473-483, aug. 2010.

LUNDGREN, J. G. **Relationships of natural enemies and non-prey foods.**

Dordrecht: Springer, 2009.

MARAZZI, B.; BRONSTEIN, J. L.; KOPTUR, S. The diversity, ecology and evolution of extrafloral nectaries: current perspectives and future challenges.

Annals of Botany, Oxford, v. 111, n. 6, p. 1243-1250, jun. 2013.

MICHAUD, J. P. Coccinellids in biological control. In: HODEK, I.; VAN EMDEN, H. F.; HONEK, A. (Ed.). **Ecology and Behaviour of the Ladybird Beetles.** Wiley-Blackwell: Oxford, 2013. p. 488-519.

MIZELL, R. F.; SCONYERS, M. C. Toxicity of imidacloprid to selected arthropod predators in the laboratory. **Florida Entomologist**, Lutz, v. 75, n. 2, p. 277-280, jun. 1992.

NAUEN, R. et al. Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. **Pesticide Biochemistry and Physiology**, San Diego, v. 76, n. 2, p. 55-69, jun. 2003.

NECHOLS, J. R.; HARVEY, T. L. Evaluation of a mechanical exclusion method to assess the impact of Russian wheat aphid natural enemies. In: QUISENBERRY, S. S.; PEAIRS, F. P. (Ed.). **Response Model for an Introduced Pest - The Russian wheat aphid.** Thomas Say Publications: Lanham, 1998. p. 270-279.

NUYTTENS, D. et al. Pesticide-laden dust emission and drift from treated seeds during seed drilling: a review. **Pest Management Science**, Hoboken, v. 69, n. 5, p. 564-575, mar. 2013.

PEMBERTON, R. W.; VANDENBERG, N. J. Extrafloral nectar feeding by ladybird beetles (Coleoptera: Coccinellidae). **Proceedings of the Entomological Society of Washington**, Washington, v. 95, n. 2, p. 139-151, 1993.

PIKE, K. S. et al. Aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) of northwest USA. **Proceedings of the Entomological Society of Washington**, Washington, v. 102, n. 3, p. 688-740, jul. 2000.

RICE, M. E.; WILDE, G. E. Experimental evaluation of predators and parasitoids in suppressing greenbugs (Homoptera: Aphididae) in sorghum and wheat. **Environmental Entomology**, Lanham, v. 17, n. 1, p. 836-841, 1988.

ROYER, T. A.; WALGENBACH, D. D. Predacious arthropods of cultivated sunflower in eastern South Dakota. **Journal of the Kansas Entomological Society**, Manhattan, v. 64, n.1, p. 112-116, jan. 1991.

SMITH, S. F.; KRISCHI, V. A. Effects of systemic imidacloprid on *Coleomegilla maculata* (Coleoptera: Coccinellidae). **Environmental Entomology**, Lanham, v. 28, n. 6, p. 1189-1195, dec. 1999.

STAPEL, J. O. et al. Extrafloral nectar, honeydew, and sucrose effects on searching behavior and efficiency of *Microplitis croceipes* (Hymenoptera: Braconidae) in cotton. **Environmental Entomology**, Lanham, v. 26, n. 3, p. 617-623, jun. 1997.

STRAUSBAUGH, C. A.; EUJAYL, I. A.; FOOTE, P. Seed treatments for the control of insects and diseases in sugarbeet. **Journal of Sugar Beet Research**, Denver, v. 47, n. 3, p. 105-125, aug. 2010.

TAYLOR, A. G.; ECKENRODE, C. J.; STRAUB, R. W. Seed coating technologies and treatments for onion: challenges and progress. **HortScience**, Alexandria, v. 36, n. 2, p. 199-205, apr. 2001.

TOMIZAWA, M.; CASIDA, J. E. Neonicotinoid insecticide toxicology: mechanisms of selective action. **Annual Review of Pharmacology and Toxicology**, Palo Alto, v. 45, n. 1, p. 247-268, feb. 2005.

WÄCKERS, F. L.; ROMEIS, J.; VAN RIJN, P. Nectar and pollen feeding by insect herbivores and implications for multitrophic interactions. **Annual Review of Entomology**, Palo Alto, v. 52, n. 1, p. 301-323, jan. 2007.

SECOND PART – ARTICLES

ARTICLE 1

**SUBLETHAL EFFECTS OF CHLORANTRANILIPROLE AND
THIAMETHOXAM SEED TREATMENTS WHEN *Lysiphlebus testaceipes*
FEED ON SUNFLOWER EXTRAFLORAL NECTAR**

This article was written in accordance with the standards of BioControl, for which it was submitted, accepted and published (BioControl: Vol. 59, No. 5, p. 503-511, 2014).

**Sublethal effects of chlorantraniliprole and thiamethoxam seed treatments
when *Lysiphlebus testaceipes* feed on sunflower extrafloral nectar**

Valéria F Moscardini,^{a,b} Pablo C Gontijo,^{a,b} JP Michaud^{b*} and Geraldo A Carvalho^a

^aDepartment of Entomology, Federal University of Lavras, Lavras, Minas Gerais, Brazil.

^bDepartment of Entomology, Kansas State University, Agricultural Research Center-Hays, Hays, Kansas, USA.

* Corresponding author: J.P. Michaud, 1232 240th Ave., Hays, KS, 67601, Email: jpmi@ksu.edu

Abstract The extrafloral nectar (EFN) of sunflower, *Helianthus annuus* L., is an important-summer resource for many insects and represents a potential route of exposure to systemic insecticides applied as seed treatments to cultivated varieties. Among the many parasitoids that utilize sunflower EFN, *Lysiphlebus testaceipes* (Cresson) (Hymenoptera: Braconidae) is an important generalist parasitoid of cereal aphids in North America. This study evaluated the performance of adult wasps fed EFN of sunflower plants grown from seed treated with chlorantraniliprole and thiamethoxam. Consumption of EFN from treated sunflower seedlings caused no lethal effects, but reduced the numbers of greenbug nymphs, *Schizaphis graminum* Rondani, attacked and parasitized when wasps foraged in Petri dish arenas. Whereas control females self-superparasitized every fourth hosts, those exposed to chlorantraniliprole did not. Offspring developmental time and adult emergence were unaffected by either treatment, but thiamethoxam greatly reduced the proportion of female offspring.

Keywords Conservation biological control Ecotoxicology *Lysiphlebus testaceipes* (Hymenoptera: Braconidae) *Schizaphis graminum* Systemic insecticides

Introduction

Plants have evolved multiple strategies to defend themselves against herbivores. Aside from direct physical and chemical defences, nectar and pollen may attract natural enemies of herbivores in addition to pollinators (Pemberton and Lee 1996; Nicolson et al. 2007). Both floral and extrafloral nectar (EFN) are an important source of hydration and nutrition for parasitoids (Heimpel and Collier 1996; Heimpel et al. 1997). The composition of EFN differs from that of floral nectar in most plant species (Baker et al. 1978). Although both are mostly sugar by weight, EFN usually contains a full complement of essential amino acids, albeit in low concentrations, with sucrose the dominant sugar, rather than glucose or fructose (Baker and Baker 1979; Rogers 1985). Sugars can be essential for parasitoid survival (Lewis et al. 1998), longevity (Dyer and Landis 1996) and fecundity (Olson and Andow 1998), and can improve parasitoid performance in biological control contexts (Gurr et al. 2004). Another benefit of EFN is that it is available to natural enemies for a much longer period than floral nectar, beginning early in plant development and continuing up to, and beyond, flowering (Pacini et al. 2003; Rose et al. 2006).

On the High Plains of the USA, sunflower plants are an important source of EFN for beneficial insects during the hot, dry summers (Royer and Walgenbach 1991). Small nectaries that occur profusely along petioles and leaf veins secrete nectar from the time the first true leaves expand until the plants senesce. A great diversity of parasitoids feed on sunflower EFN (e.g., Charlet and Gavloski 2011) and the presence of sunflowers has been shown to improve

the area-wide survival of natural enemy species important in the biological control of key pests in neighboring crops (e.g., Brewer et al. 2008), probably by supplying critical resources for natural enemies (Gurr et al. 2004). Intercropping cultivated plants with EFN producing companion plants has even been proposed to supplementary food for natural enemies early in the growing season before pests become abundant (Jamont et al. 2013).

Like many other row crops cultivated on the High Plains, the majority of commercial sunflower seed is now planted with a systemic insecticide seed treatment which poses a potential hazard to natural enemies. Seed treatments have been promoted as more compatible with biological control than broadcast sprays of insecticides, primarily because seed treatment involves a smaller amount of pesticide applied in a more selective manner, with consequently reduced impacts on non-target organisms (Hull and Beers 1985; Albajes et al. 2003) and lower levels of environmental contamination (Taylor et al. 2001). Systemic insecticides typically show low lipophilicity and exhibit a low octanol-water partition coefficient ($\log P_{oct}$) (Cloyd and Bethke 2011). This property facilitates their translocation in plant tissues, and may lead to contamination of pollen, floral and EFN (Maienfisch et al. 2001; Lahm et al. 2009) that, in turn, can cause both lethal and sublethal effects on natural enemies over potentially prolonged periods (Lundgren 2009; Li et al. 2012). EFN is produced in much earlier stages of plant development than floral nectar and thus has the potential to bear higher concentrations of systemic insecticides applied as seed treatments. For example, residues of the neonicotinoid imidacloprid were 50 times more concentrated in cotyledons than in the first true leaves of sunflower, and 800 times more concentrated than in the apex of plant. Concentrations were typically reduced by half in each successive pair of leaves, reflecting progressive dilution as a function of plant growth (Laurent and Rathahao 2003).

Chlorantraniliprole and thiamethoxam represent two different systemic insecticide groups used in seed treatment, diamides and neonicotinoids, respectively. Diamide insecticides have a novel mode of action that acts exclusively on the ryanodine receptor in insect neuromuscular junctions (Nauen 2006; Lahm et al. 2009). In contrast, thiamethoxam is a neonicotinoid insecticide that targets nicotinic acetylcholine receptors in the insect central nervous system (Tomizawa and Casida 2005) and can be converted within the plant into highly toxic metabolites such as clothianidin (Casida 2011) which can increase its toxicity.

Lysiphlebus testaceipes (Cresson) (Hymenoptera: Braconidae) is a solitary endoparasitoid that attacks more than 100 aphid species, mostly on grasses and herbaceous plants (Pike et al. 2000). It is an important natural enemy of the greenbug, *Schizaphis graminum* (Rondani) (Hemiptera: Aphididae) and helps to maintain populations of this aphid below economic levels in both sorghum (Jones 2001) and wheat (Giles et al. 2003). Without nectar, many parasitoids show a dramatic reduction in their ability to parasitize insect pests (Stapel et al. 1997; Lewis et al. 1998), and adult *L. testaceipes* can often be observed feeding on sunflower EFN in the High Plains environment (J.P.M. unpublished observations). Therefore, the objectives of this study were to assess whether either chlorantraniliprole or thiamethoxam would have any impacts on the survival of adult wasps, their foraging behavior, or the development of their offspring, when the adults fed on EFN of sunflower seedlings grown from treated seed.

Materials and methods

Insect colonies

All insect colonies were held in growth chambers under the same physical conditions: $21 \pm 1^\circ\text{C}$ and a L:D 16:8 photoperiod. The colony of *S. graminum* was initiated from aphids naturally infesting sorghum in a greenhouse at the Agricultural Research Center-Hays, in Hays, KS, USA. The aphids were established on seedlings of a greenbug-susceptible sorghum variety (cv. P8500) planted in metal trays (8.0 cm x 51.0 cm x 36.0 cm) filled with a mixture of soil, peat moss and perlite (1:1:1) and germinated in a greenhouse at $25 \pm 2^\circ\text{C}$. A colony of *L. testaceipes* was established from material obtained from a laboratory culture that had been maintained on *Rhopalosiphum padi* (L.) (Hemiptera: Aphididae) at the Department of Entomology, University of Minnesota, MN, USA for many generations. The colony was reared on *S. graminum* for three generations prior to use in bioassays. For each generation, mated parasitoid females (2-3 days old, $n = 20$) were each provided ca. 50 2nd and 3rd instar aphids in a Petri dish (5.5 cm diam) for a period of 50 min tallied from the time the first aphid was stung. Following removal of wasps, the aphids were transferred to sorghum seedlings in plastic pots (15.0 cm diam x 14.0 cm ht), ca. 250 aphids per pot. The seedlings had been germinated in a mixture of soil, peat moss and perlite (1:1:1) in a greenhouse at $25 \pm 2^\circ\text{C}$. The pots were placed in a growth chamber and, once mummification of parasitized aphids occurred, mummies were removed from the plants and placed in a wax paper cup (8.5 cm diam x 12.5 ht) covered with a plastic Petri lid until adult emergence. Emergent adults were fed with a solution of diluted honey (50 %) streaked onto the plastic lid. Males and females were held together for ca. 48 h to permit mating before females were removed and used to produce the next generation.

Sunflower plants

Seeds of Pioneer 63N82 sunflower treated with chlorantraniliprole (1,800 mg a.i. 100 kg⁻¹) were provided by DuPont Crop Protection (division of E.I. du Pont de Nemours and Co., Wilmington, DE, USA). Seeds of Triumph Nusun cv. 810CL were obtained from Triumph Seed Corp. (division of Dow Agrosiences, Indianapolis, ID, USA) both with and without treatment with Cruiser 5FS[®] (thiamethoxam, 50 mg a.i. 100 kg⁻¹, Syngenta Crop Protection, Greensboro, NC, USA). Untreated Triumph seed served as the experimental control. Seeds for each treatment were planted in separate metal trays (8.0 cm x 51.0 cm x 36.0 cm) filled with a mixture of soil, peat moss and perlite (1:1:1) and germinated in a greenhouse at 25 ± 2 °C under natural light supplemented during daylight hours with metal halide lamps (L:D =12:12). Plants were watered daily, but sparingly, to avoid excessive leaching of insecticide. Sunflower stalks were harvested beginning at the V2 stage (two true leaves expanded) and every two days thereafter throughout each period of parasitoid exposure in experimental treatments. This growth stage corresponded to 14-15 days-old plants, post-emergence, under the prevailing temperature conditions. Sunflower plants begin secreting extrafloral nectar shortly about two weeks after germination, the first nectaries developing on the main stem between the cotyledons and the first true leaves. These small nectaries are invisible to the naked eye, but they occur profusely over the surface of stems, leaf petioles and leaf veins. The nectar is exuded in very small amounts from these tiny pores and it appears to require considerable feeding effort on the part of insects to obtain the EFN. For provisioning to insects, stem segments (ca. 10.0 cm long) were excised from seedlings and the cut ends dipped in liquid paraffin to seal vascular tissues and maintain turgor, while at the same time preventing the exudation of any resinous materials that might pose a hazard to the insects.

Experimental procedure

In order to test for lethal and sublethal effects of exposure to EFN from treated and untreated seedlings, mummies of *L. testaceipes* (n = 30) were placed in wax paper cups (n = 4 cups per treatment), each containing two sunflower stem segments and sealed with a plastic Petri dish lid. Upon emergence, the adult wasps were exposed to the sunflower stems as their only food source for a period of 48 h, with all stems replaced after 24 h. Mortality was then tallied after 48 h with cup as the experimental unit. Sublethal effects on behavior were assessed by aspirating individual females (n = 16 per treatment) into Petri dishes (5.5 cm diam), each containing 40 2nd and 3rd instar aphids. Following release of each female wasp, the time to first sting (attack latency) was recorded and the wasp was left to forage for 30 min, whereupon the female was removed. All aphids from each dish were then transferred with a fine brush to a single sorghum seedling (one per replicate) that had been germinated in a 16.0 cm plastic cone (Stuewe & Sons, Corvallis, OR, USA) filled with a mixture of soil, peat moss and perlite (1:1:1). Wasps that did not sting an aphid within 10 min of release were replaced. Each cone was then covered with a custom-made, clear plastic cylinder (30 cm length), sealed at the top with a plastic plug and ventilated on the sides by means of a series of screened holes. The cones were inserted into a supporting rack and transferred to a growth chamber. Plants were watered every 48 h by submerging the rack of cones in a water bath for 30 min. Four days after parasitism, a sample of aphids from each plant (n = 10) were dissected in saline under a stereo microscope (40x magnification) to verify the presence/absence of parasitoid larvae. The attack rate was estimated as the sum of parasitized aphids (those containing a parasitoid larvae) and pseudoparasitized aphids (those with teratocytes present and evident deterioration of aphid embryos, but no parasitoid larva). Aphids containing more

than one wasp larva were tallied as self-superparasitized. As mummies formed, they were carefully removed from plants and transferred to a wax paper cup (as above) until emergence of adults.

Data analysis

Data were subjected to one-way ANOVA and, when significant, means were separated using Fisher's LSD test ($\alpha = 0.05$) (PROC GLM; SAS Institute 2008). Exposure cups were considered replicates for analysis of mortality during exposure, individual females as replicates for analysis of attack behavior, and sorghum seedlings each bearing the aphids exposed to a single female as replicates for analysis of progeny fitness. Attack latency data were $\log(x + 1)$ transformed, whereas percentage data (percent hosts attacked, parasitized, and offspring emergence) were arcsine square root transformed to fit requirements for normality and homoscedasticity (PROC UNIVARIATE; SAS Institute 2008). Untransformed means are presented in figures and tables. Proportional data (self-superparasitism, mummification, and sex ratio ($\Sigma \text{♀} / \Sigma (\text{♀} + \text{♂})$)) were analyzed using the χ^2 test ($\alpha = 0.05$) (PROC FREQ; SAS Institute 2008).

Results

Neither chlorantraniliprole nor thiamethoxam seed treatments produced mortality of *L. testaceipes* any higher than controls after 48 h when adults consumed the EFN of excised sunflower stem segments (percent mortality \pm SE: control = 20.8 ± 5.0 ; chlorantraniliprole = 18.3 ± 7.5 ; thiamethoxam = 19.2 ± 6.4 ; $F_{2,9} = 0.04$, $P = 0.961$). However, both insecticide treatments reduced the latent period to first sting, number of aphids stung, and number of aphids

parasitized (Table1). Self-superparasitism was reduced by the chlorantraniliprole treatment, indicating lower levels of oviposition, and thiamethoxam reduced the offspring sex ratio (proportion female). Neither the proportion of remaining aphids forming mummies, adult emergence, nor development time from parasitism to emergence, were significantly affected by either treatment (Table2).

Discussion

Although no direct wasp mortality was observed in this study, the consumption of EFN from sunflower seedlings treated with chlorantraniliprole and thiamethoxam caused various sublethal effects on *L. testaceipes* behavior and biological performance. Female wasps exposed to either material were faster to attack their first aphid compared to control females, although they attacked fewer aphids within the test period. These females displayed various combinations of erratic behavior, disorientated movements, and extended periods of antennation and ovipositor probing on the same host, suggesting impaired host acceptance behavior. We suspect that the time to first host encounter was faster for treated females because they moved randomly within the dish from time of introduction, whereas control wasps spent an initial period evaluating sensory cues associated with their surrounding environment prior to initiating search behavior. For example, Hopkinson et al. (2013) observed that female *L. testaceipes* foraging for *Aphis gossypii* Glover and *Aphis craccivora* (Koch) (Hemiptera: Aphididae) on cotton leaf discs required a mean of 14.7 and 16.4 min, respectively, to attack their first host, illustrating how much time may be expended by these insects exploring a leaf surface prior to an initial host encounter. Therefore, it is reasonable to expect that other behavioral impairments might be evident in a more complex, natural environment, that

would not be observed in the experimental arenas. The relatively short latent periods observed in the present study are likely the result of presenting hosts in a small arena in the absence of plant material or other cues such as honeydew. Residues of various insecticides are known to trigger changes in parasitoid mobility, orientation, feeding and parasitism ability (Desneux et al. 2007; Garcia 2011). Changes in foraging behavior and rates of parasitism were observed after *Anagrus nilaparvatae* (Pang and Wang) (Hymenoptera: Mymaridae) fed on a mixture of honey and imidacloprid (Liu et al. 2010) and subsequent work found similar effects of chlorantraniliprole on the same species (Liu et al. 2012). Both a braconid and an encyrtid parasitoid that fed on floral nectar of *Eucalyptus* trees treated with an imidacloprid soil drench suffered reduced survival and reproductive performance (Paine et al. 2011), although in this case the concentrations detected in nectar were more than double the LC50 values for these species. In a three year field study, cotton seeds treated with either imidacloprid or thiamethoxam reduced the season-long rate of whitefly, *Bemisia tabaci* (Gennadius) (Hemiptera: Aleyrodidae), parasitism by a complex of three aphelinid parasitoids (Naveed et al. 2010).

The similar reductions in parasitoid performance caused by the chlorantraniliprole and thiamethoxam treatments likely result from entirely different effects on the insects, as the former material interferes with muscle contraction, and the latter with neurotransmission. Chlorantraniliprole induces the ryanodine receptors to release stored calcium ions from the sarcoplasmic reticulum, causing impaired regulation of muscle contraction, paralysis and ultimately death in sensitive species (Cordova et al. 2006; Lahm et al. 2007). In contrast, thiamethoxam targets nicotinic acetylcholine receptors on the post-synaptic membranes of nerve cell junctions producing both lethal and sublethal neurological effects (Tomizawa and Casida 2005). Both materials appeared to lower the overall mobility of females, resulting in fewer host encounters during

the test period, and consequently less parasitism. This can be construed to result entirely from impaired muscle function in the chlorantraniliprole treatment. However, the neurotoxic activity of thiamethoxam has the potential to both impede wasp motor functions and their sensory perception of cues involved in host evaluation and acceptance. Disruption of parasitoid sensory perception and motor function as a function of sublethal exposure to neurotoxic insecticides is well documented (Haynes 1988; Garcia 2011). For example, Komeza et al. (2001) studied *Leptopilina boulardi* (Hymenoptera: Figitidae) females exposed to the LD20 of the organophosphate chlorpyrifos and found them to be less efficient in finding host patches marked with kairomones, and exhibit longer patch residence times. In contrast, Delpuech et al. (2005) found that the congeneric *Leptopilina heterotoma* Thomson females responded to kairomones of *Drosophila* with greater arrestment following contact with chlorpyrifos and deltamethrin at LD20. Thus, either inhibitive or stimulative (hormoligosis) effects of neurotoxic insecticides may be observed at sublethal doses, depending on the insecticide and species of insect.

Self-superparasitism occurs when a female lays eggs in a host that she has previously parasitized (Waage 1986). In contrast to control females that laid an additional egg in almost every fourth host, *L. testaceipes* females exposed to chlorantraniliprole did not engage in this behavior (Table 1). Although *L. testaceipes* is a solitary parasitoid and only one offspring can survive per host, there exist several potential payoffs for self-superparasitism. Aphid parasitoids such as *L. testaceipes* emerge with hundreds of eggs ready to lay and laying an additional egg in some hosts can improve a female's probability of host possession for her offspring if there is a risk of superparasitism by conspecific females (Michaud and Mackauer 1995). Although supernumerary larvae are eliminated, upon hatching, additional eggs release additional teratocytes critical to selective digestion of host tissues (Falabella et al. 2000; Caccia et al. 2012)

and these may improve the survival of neonate parasitoids in their physiological struggle against host resistance mechanisms. For example, Bai and Mackauer (1992) found that *Aphidius ervi* Halliday developing in superparasitized pea aphids, *Acyrtosiphum pisum* Harris, gained 14 % more dry mass compared to those in singly-parasitized hosts, with no cost in additional developmental time. Similarly, Zhang et al. (2010) reported that oviposition of multiple eggs increased parasitism success of the solitary parasitoid *Campoletis chloridae* Uchida (Hymenoptera: Ichneumonidae) in larvae of *Mythimna separate* Walker (Noctuidae: Lepidoptera). In keeping with the inference that self-superparasitism of some fraction of hosts is an adaptive 'insurance' strategy for solitary parasitoids, its absence in females exposed to chlorantraniliprole can be considered an abnormality likely to have adverse effects on female fitness, for example if the behavior is adaptive for improving offspring survival in hosts of marginal suitability.

The lack of any significant difference among treatments in the proportion of aphids mummified appears somewhat inconsistent with the results obtained from dissected subsamples, and likely reflects a bias toward sampling parasitized aphids during removal of these subsamples. In order to minimize disturbance of the remaining aphids, subsamples were removed from sorghum seedlings by gently tapping them over a sheet of paper and collecting the first ten aphids so dislodged. Many factors can potentially influence the dropping propensity of aphids from a plant, including parasitism (e.g. McAllister and Roitberg 1987). Thus, we suspect that parasitized aphids had greater dropping propensity than unparasitized aphids and were thus over-represented in the dissected samples and under-represented in the remaining fraction that was reared to mummification.

Neither offspring survival nor their developmental time was affected by either seed treatment (Table 2), but the sex ratio (proportion female) was

significantly reduced by the thiamethoxam treatment. In arrhenotokous Hymenoptera such as *L. testaceipes*, unfertilized eggs become males, so egg fertilization (and thus sex ratio) is actively controlled, and females may even selectively place fertilized eggs in better hosts (e.g. Srivastava and Singh 1995a). In aphidiine wasps, sex ratio can be diminished by various stressors such as high temperature (Deng and Tsai 1998; Matin et al. 2009), low temperature (Sigsgaard 2000; Mahi et al. 2014), multiple matings (Kant et al. 2012) and aging (Srivastava and Singh 1995b; Kant et al. 2013). Since the rate of mummification among aphids not harvested for dissection was similar across treatments, the lower sex ratio in the thiamethoxam treatment is most likely attributable to neurologic impairment of females that interfered to some extent with their ability to fertilize eggs. Since both parasitoid population growth and biological control efficacy hinge on numbers of parasitoid females, a reduction in sex ratio to only 11 % female can be considered a highly adverse result.

In summary, many studies have emphasized that systemic insecticides used in seed treatments pose risks to beneficial species such as parasitoids (Prabhaker et al. 2011), predators (Al-Deeb et al. 2001; Moser and Obrycki 2009), and pollinators (Girolami et al. 2009) due to their translocation within plants and their potential contamination of pollen and nectar (Cloyd and Bethke 2011). The treatment of seed with such insecticides is now a widespread practice in production agriculture because it can improve stand establishment in row crops by protecting germinating seeds and young seedlings. However, neonicotinoids are notoriously incompatible with integrated pest management (IPM) (Hutchins 2010), can accumulate in soils, and are prone to leaching into waterways (Goulson 2013). Furthermore, there are risks to pollinators associated with improper handling or disposal of residues during planting operations (Nuyttens et al. 2013). Work in soybean has questioned whether producers gain any significant benefits from seed treatments (Seagraves and Lundgren 2012).

Producers buying factory-treated seed incur a control cost prior to suffering any pest infestation, an expense that is not recouped if economically damaging pests do not attack plants in early growth stages. Because seed treatments constitute a prophylactic control, they are not consistent with IPM principles that require assessment of pest numbers and projection of an economic impact prior to resorting to pesticide application (Stern et al. 1959). Further studies are warranted to determine just how compatible seed treatments are with IPM programs that rely on conservation biological control to any significant extent.

Acknowledgements The authors are grateful to the CAPES Foundation (Brazilian Ministry of Education), the National Council of Scientific and Technological Development (CNPq), and the Minas Gerais State Foundation for Research Aid (FAPEMIG) for scholarship support from CAPES - no. 3362-13-2 (VFM) and CAPES - no. 3363-13-9 (PCG). This is contribution No. 14-325-J of the Kansas State Experiment Station.

References

- Albajes R, López C, Pons X (2003) Predatory fauna in cornfields and response to imidacloprid seed treatment. *J Econ Entomol* 96:1805-1813
- Al-Deeb MA, Wilde GE, Zhu KY (2001) Effect of insecticides used in corn, sorghum, and alfalfa on the predator *Orius insidiosus* (Hemiptera: Anthocoridae). *J Econ Entomol* 94:1353-1360
- Bai B, Mackauer M (1992) Influence of superparasitism on development rate and adult size in a solitary parasitoid wasp, *Aphidius ervi*. *Func Ecol* 6:302-307
- Baker I, Baker HG (1979) Chemical constituents of the nectars of two *Erythrina* species and their hybrid. *Ann Mo Bot Gard* 66:446-450

- Baker HG, Opler PA, Baker I (1978) A comparison of amino-acid complements of floral and extrafloral nectars. *Bot Gaz* 139:322-332
- Brewer MJ, Noma T, Elliott NC, Kravchenko AN, Hild AL (2008) A landscape view of cereal aphid parasitoid dynamics reveals sensitivity to farm- and region-scale vegetation structure. *Eur J Entomol* 105:503-511
- Caccia S, Grimaldi A, Casartelli M, Falabella P, Eguileor M de, Pennacchio F, Giordana B (2012) Functional analysis of a fatty acid binding protein produced by *Aphidius ervi* teratocytes. *J Insect Physiol* 58:621-627
- Casida JE (2011) Neonicotinoid metabolism: compounds, substituents, pathways, enzymes, organisms, and relevance. *J Agric Food Chem* 59:2923-2931
- Charlet LD, Gavloski J (2011) Insects of sunflower in the Northern Great Plains of North America. In: Floate KD (ed) *Arthropods of canadian grasslands: inhabitants of a changing landscape*, vol 2. Biological Survey of Canada, pp 159-178
- Cloyd RA, Bethke JA (2011) Impact of neonicotinoid insecticides on natural enemies in greenhouse and interiorscape environments. *Pest Manag Sci* 67:3-9
- Cordova D, Benner EA, Sacher MD, Rauh JJ, Sopa JS, Lahm GP, Selby TP, Stevenson TM, Flexner L, Gutteridge S, Rhoades DF, Wu L, Smith RM, Tao Y (2006) Anthranilic diamides: A new class of insecticides with a novel mode of action, ryanodine receptor activation. *Pestic Biochem Phys* 84:196-214
- Delpuech JM, Bardon C, Bouletreau M (2005) Increase of the behavioral response to kairomones by the parasitoid wasp *Leptopilina heterotoma* surviving insecticides. *Arch Environ Con Tox* 49:186-191

- Deng YX, Tsai JH (1998) Development of *Lysiphlebia japonica* (Hymenoptera: Aphidiidae), a parasitoid of *Toxoptera citricida* (Homoptera: Aphididae) at five temperatures. Fla Entomol 81:415-423
- Desneux N, Decourtye A, Delpuech JM (2007) The sublethal effects of pesticides on beneficial arthropods. Annu Rev Entomol 52:81-106
- Dyer LE, Landis DA (1996) Effects of habitat, temperature, and sugar availability on longevity of *Eriborus terebrans* (Hymenoptera: Ichneumonidae). Environ Entomol 25:1192-1201
- Falabella P, Tremblay E, Pennacchio F (2000) Host regulation by the aphid parasitoid *Aphidius ervi*: the role of teratocytes. Entomol Exp Appl 97:1-9
- Garcia P (2011) Sublethal effects of pyrethroids on insect parasitoids: what we need to further know. In: Stoytcheva M (ed) Pesticides - formulations, effects, fate. InTech, Rijeka, Croatia, pp 477-494
- Giles KL, Jones DB, Royer TA, Elliott NC, Kindler SD (2003) Development of a sampling plan in winter wheat that estimates cereal aphid parasitism levels and predicts population suppression. J Econ Entomol 96:975-982
- Girolami V, Mazzon L, Squartini A, Mori N, Marzaro M, Di bernardo A, Greatti M, Giorio C, Tapparo A (2009) Translocation of neonicotinoid insecticides from coated seeds to seedling guttation drops: a novel way of intoxication for bees. J Econ Entomol 102:1808-1815
- Goulson D (2013) REVIEW: An overview of the environmental risks posed by neonicotinoid insecticides. J Appl Ecol 50:977-987
- Gurr GM, Scarratt SL, Wratten SD, Berndt L, Irvin NA (2004) Ecological engineering, habitat manipulation and pest management. In: Gurr GM, Wratten SD, Altieri MA (eds) Ecological Engineering for Pest Management. Comstock Press, Ithaca, USA, pp 1-12

- Haynes KF (1988) Sublethal effects of neurotoxic insecticides on insect behavior. *Annu Rev Entomol* 33:149-168
- Heimpel GE, Collier TR (1996) The evolution of host-feeding behaviour in insect parasitoids. *Biol Rev* 71:373-400
- Heimpel GE, Rosenheim JA, Kattari D (1997) Adult feeding and lifetime reproductive success in the parasitoid *Aphytis melinus*. *Entomol Exp Appl* 83:305-315
- Hopkinson JE, Zalucki MP, Murray DAH (2013) Host selection and parasitism behavior of *Lysiphlebus testaceipes*: role of plant, aphid species and instar. *Biol Control* 64:283-290
- Hull L, Beers E (1985) Ecological selectivity: modifying chemical control practices to preserve natural enemies. In: Hoy MA, Herzog DC (eds) *Biological control in agricultural IPM systems*. Academic Press, New York, USA, pp. 103-122
- Hutchins SH (2010) Indifference analysis: a practical method to assess uncertainty in IPM decision making. *J Integ Pest Manag* 1:D1-D3
- Jamont M, Crépeillère S, Jaloux B (2013) Effect of extrafloral nectar provisioning on the performance of the adult parasitoid *Diaeretiella rapae*. *Biol Control* 65:271-277
- Jones DB (2001) Natural enemy thresholds for greenbug, *Schizaphis graminum* Rondani, on winter wheat. Oklahoma State University, Stillwater, USA
- Kant R, Trewick SA, Sandanayaka WRM, Godfrey AJR, Minor MA (2012) Effects of multiple matings on reproductive fitness of male and female *Diaeretiella rapae*. *Entomol Exp Appl* 145: 215-221
- Kant R, Minor MA, Sandanayaka WRM, Trewick SA (2013) Effects of mating and oviposition delay on parasitism rate and sex allocation behaviour of *Diaeretiella rapae* (Hymenoptera: Aphidiidae). *Biol Control* 65:265-270

- Komezha N, Fouillet P, Bouletreau M, Delpuech JM (2001) Modification, by the insecticide chlorpyrifos, of the behavioral response to kairomones of a parasitoid wasp, *Leptopilina boulandi*. Arch Environ Con Tox 41:436-442
- Lahm GP, Stevenson TM, Selby TP, Freudenberger JH, Cordova D, Flexner L, Bellin CA, Dubas CM, Smith BK, Hughes KA, Hollingshaus JG, Clark CE, Benner EA (2007) Rynaxypyr: a new insecticidal anthranilic diamide that acts as a potent and selective ryanodine receptor activator. Bioorg Med Chem Lett 17:6274-6279
- Lahm GP, Cordova D, Barry JD (2009) New and selective ryanodine receptor activators for insect control. Bioorg Med Chem 17:4127-4133
- Laurent FM, Rathahao E (2003) Distribution of [¹⁴C] imidacloprid in sunflowers (*Helianthus annuus* L.) following seed treatment. J Agric Food Chem 51:8005-8010
- Lewis WJ, Stapel JO, Cortesero AM, Takasu K (1998) Understanding how parasitoids balance food and host needs: importance to biological control. Biol Control 11:175-183
- Li X, Degain BA, Harpold VS, Marcon PG, Nichols RL, Fournier AJ, Naranjo SE, Palumbo JC, Ellsworth PC (2012) Baseline susceptibilities of B- and Q-biotype *Bemisia tabaci* to anthranilic diamides in Arizona. Pest Manag Sci 68:83-91
- Liu F, Bao SW, Song Y, Lu HY, Xu JX (2010) Effects of imidacloprid on the orientation behavior and parasitizing capacity of *Anagrus nilaparvatae*, an egg parasitoid of *Nilaparvata lugens*. BioControl 55:473-483
- Liu F, Zhang X, Gui Q-Q, Xu Q-J (2012) Sublethal effects of four insecticides on *Anagrus nilaparvatae* (Hymenoptera: Mymaridae), an important egg parasitoid of the rice planthopper *Nilaparvata lugens* (Homoptera: Delphacidae). Crop Prot 37:13-19

- Lundgren JG (2009) Relationships of natural enemies and non-prey foods. Springer, Dordrecht, Netherlands
- Mahi H, Rasekh A, Shishehbor P, Michaud JP (2014) The biology of *Lysiphlebus fabarum* (Braconidae, Aphidiinae) following cold storage of larvae and pupae under various thermal regimes. Entomol Exp Appl (in press).
- Maienfisch P, Angst M, Brandl F, Fischer W, Hofer D, Kayser H, Kobel W, Rindlisbacher A, Senn R, Steinemann A, Widmer Hr (2001) Chemistry and biology of thiamethoxam: a second generation neonicotinoid. Pest Manag Sci 57:901-913
- Matin SB, Sahragard A, Rasoolian G (2009) Some biological parameters of *Lysiphlebus fabarum* (Hymenoptera: Aphidiidae) a parasitoid of *Aphis fabae* (Homoptera: Aphidiidae) under laboratory conditions. Munis Entomol Zool 4:193-200
- McAllister MK, Roitberg BD (1987) Adaptive suicidal behavior in pea aphids. Nature 328:797-799
- Michaud JP, Mackauer M (1995) Oviposition behavior of *Monoctonus paulensis* (Hymenoptera: Aphidiidae): factors influencing reproductive allocation to hosts and host patches. Ann Entomol Soc Am 88:220-226
- Moser SE, Obrycki JJ (2009) Non-target effects of neonicotinoid seed treatments; mortality of coccinellid larvae related to zoophytophagy. Biol Control 51:487-492
- Nauen R (2006) Insecticide mode of action: return of the ryanodine receptor. Pest Manag Sci 68:690-692
- Naveed M, Salam A, Saleem MA, Rafiq M, Hamza A (2010) Toxicity of thiamethoxam and imidacloprid as seed treatments to parasitoids associated to control *Bemisia tabaci*. Pak J Zool 42:559-565

- Nicolson S, Nepi I, Pacini E (2007) Nectaries and Nectar. Springer, Heidelberg, Germany
- Nuyttens D, Devarrewaere W, Verboven P, Foque D (2013) Pesticide-laden dust emission and drift from treated seeds during seed drilling: a review. *Pest Manag Sci* 69:564-575
- Olson DM, Andow DA (1998) Larval crowding and adult nutrition effects on longevity and fecundity of female *Trichogramma nubilale* Ertle & Davis (Hymenoptera: Trichogrammatidae). *Environ Entomol* 27:507-512
- Pacini E, Nepi M, Vesprini JL (2003) Nectar biodiversity: a short review. *Plant Syst Evol* 238:7-21
- Paine TD, Hanlon CC, Byrne FJ (2011) Potential risks of systemic imidacloprid to parasitoid natural enemies of a cerambycid attacking *Eucalyptus*. *Biol Control* 56:175-178
- Pemberton RW, Lee JH (1996) The influence of extrafloral nectaries on parasitism of an insect herbivore. *Am J Bot* 83:1187-1194
- Pike KS, Stary P, Miller T, Graf G, Allison D, Boyd-ston L, Miller R (2000) Aphid parasitoids (Hymenoptera: Braconidae: Aphidiinae) of northwest USA. *Proc Entomol Soc Wash* 102:688-740
- Prabhaker N, Castle SJ, Naranjo SE, Toscano NC, Morse JG (2011) Compatibility of two systemic neonicotinoids, imidacloprid and thiamethoxam, with various natural enemies of agricultural pests. *J Econ Entomol* 104:773-781
- Rogers CE (1985) Extrafloral nectar: ecological implications. *Bull Entomol Soc Am* 31:15-20
- Rose USR, Lewis J, Tumlinson JH (2006) Extrafloral nectar from cotton (*Gossypium hirsutum*) as a food source for parasitic wasps. *Funct Ecol* 20:67-74

- Royer TA, Walgenbach DD (1991) Predacious arthropods of cultivated sunflower in Eastern South Dakota. *J Kansas Entomol Soc* 64:112-116
- SAS Institute (2008) SAS for Windows Version 9.0. SAS Institute, Cary, USA
- Seagraves MP, Lundgren JG (2012) Effects of neonicotinoid seed treatments on soybean aphid and its natural enemies. *J Pest Sci* 85:125-132
- Sigsgaard S. (2000) The temperature-dependent duration of development and parasitism of three cereal aphid parasitoids, *Aphidius ervi*, *A. rhopalosiphi*, and *Praon volucre*. *Entomol Exp Appl* 95:173-184
- Srivastava M, Singh R (1995a) Sex ratio adjustment by a koinobiotic parasitoid *Lysiphlebus delhiensis* (Subba Rao & Sharma) (Hymenoptera: Aphidiidae) in response to host size. *Biol Agric Hort* 12:15-28
- Srivastava M, Singh R (1995b) Influence of age of parents *Lysiphlebus delhiensis* (Subba Rao and Sharma) (Hym., Aphidiidae) during copulation on progeny production and offspring sex ratio. *J Appl Entomol* 119:72-77
- Stapel JO, Cortesero AM, De Moraes CM, Tumlinson JH, Lewis WJ (1997) Extrafloral nectar, honeydew, and sucrose effects on searching behavior and efficiency of *Microplitis croceipes* (Hymenoptera: Braconidae) in cotton. *Environ Entomol* 26:617-623
- Stern VM, Smith RF, van den Bosch R, Hagen KS (1959) The Integrated Control Concept. *Hilgardia* 29:81-101
- Taylor AG, Eckenrode CJ, Straub RW (2001) Seed coating technologies and treatments for onion: Challenges and progress. *HortScience* 36:199-205
- Tomizawa M, Casida JE (2005) Neonicotinoid insecticide toxicology: mechanisms of selective action. *Annu Rev Pharmacol* 45:247-268
- Waage JK (1986) Family planning in parasitoids: adaptative patterns of progeny and sex allocation. In: Waage JK, Greathead D (eds) *Insect Parasitoids*. Academic Press, London, UK, pp 63-95

Zhang JH, Gu LQ, Wang CZ (2010) Superparasitism behavior and host discrimination of *Campoletis chlorideae* (Ichneumonidae: Hymenoptera) toward *Mythimna separata* (Noctuidae: Lepidoptera). *Environ Entomol* 39:1249-1254

Table 1 Mean (\pm SE) parameters of *Lysiphlebus testaceipes* parasitism behavior when females were subjected to three treatments (wasps fed for 48h on extrafloral nectar of sunflower stems excised from seedlings grown from seed treated with either chlorantraniliprole or thiamethoxam, or controls) prior to foraging as individuals on 40 second and third instar nymphs of *Schizaphis graminum* in a plastic Petri dish for 30 min.

Seed treatment	Attack latency ^a (min)	Percent attacked ^a	Percent parasitized ^a	Self-superparasitism ^{b,c}
Control	3.0 \pm 0.7 a	43.1 \pm 7.4 a	38.8 \pm 7.9 a	0.24 a
Chlorantraniliprole	1.5 \pm 0.4 b	21.9 \pm 6.0 b	19.4 \pm 5.7 b	0.00 b
Thiamethoxam	1.1 \pm 0.3 b	24.4 \pm 5.2 b	20.6 \pm 5.0 b	0.09 ab
<i>F</i> or χ^2	4.55	4.39	3.56	10.86
<i>df</i>	2,45	2,45	2,45	2
<i>P</i>	0.016	0.018	0.037	0.004

Ten aphids were dissected from each replicate to determine percentages attacked and parasitized, and the proportion self-superparasitized.

Means (\pm SE) and proportions followed by different letters were significantly different within columns (Fisher's LSD, or $\chi^2 \alpha = 0.05$).

^a Analysis by one-way ANOVA, or ^b χ^2

^c Proportion of dissected hosts that contained two larvae.

Table 2 Mean (\pm SE) developmental parameters of *Lysiphlebus testaceipes* progeny whose mothers were subjected to three treatments (wasps fed for 48h on extrafloral nectar of sunflower stems excised from seedlings grown from seed treated with chlorantraniliprole or thiamethoxam, or controls) prior to parasitizing second and third instar nymphs of *Schizaphis graminum* in a plastic Petri dish.

Seed treatment	Proportion aphids Mummified ^b	Adult emergence ^a (%)	Development time ^a (days)	Sex ratio ^b (proportion female)
Control	0.25 \pm 0.02 a	82.2 \pm 5.2 a	15.7 \pm 0.2 a	0.41 \pm 0.05 a
Chlorantraniliprole	0.19 \pm 0.02 a	79.3 \pm 7.9 a	15.8 \pm 0.2 a	0.40 \pm 0.06 a
Thiamethoxam	0.21 \pm 0.02 a	72.6 \pm 4.2 a	15.5 \pm 0.1 a	0.11 \pm 0.04 b
<i>F</i> or χ^2	4.19	0.71	0.74	20.15
<i>df</i>	2	2,35	2,35	2
<i>P</i>	0.123	0.498	0.485	< 0.001

Values followed by different letters were significantly different within columns (χ^2 , $\alpha = 0.05$).

^a Data were analyzed by one-way ANOVA, or ^b χ^2 .

ARTICLE 2

**SUBLETHAL EFFECTS OF INSECTICIDE SEED TREATMENTS ON
TWO NEARCTIC LADY BEETLES (COLEOPTERA:
COCCINELLIDAE)**

This article was written in accordance with the standards of Ecotoxicology, for
which it was submitted.

**Sublethal effects of insecticide seed treatments on two nearctic lady beetles
(Coleoptera: Coccinellidae)**

Running head: Toxicity of seed treatments to lady beetles

Valéria Fonseca Moscardini,^{a,b} Pablo Costa Gontijo,^{a,b} JP Michaud^{b*} and
Geraldo Andrade Carvalho^a

^aDepartment of Entomology, Federal University of Lavras, Lavras, Minas Gerais, Brazil.

^bDepartment of Entomology, Kansas State University, Agricultural Research Center-Hays, Hays, Kansas, USA.

*Corresponding author: J.P. Michaud, 1232 240th Ave., Hays, KS, 67601,
Email: jpmi@ksu.edu

Abstract Predatory insects often feed on plants or use plant products to supplement their diet, creating a potential route of exposure to systemic insecticides used as seed treatments. This study examined whether chlorantraniliprole or thiamethoxam might negatively impact *Coleomegilla maculata* and *Hippodamia convergens* when the beetles consumed the extrafloral nectar of sunflowers grown from treated seed. We reared both species on eggs of *Ephestia kuehniella* and then switched adult *H. convergens* to a diet of greenbugs, *Schizaphis graminum*, in order to induce oviposition in this species. Excised sunflower stems, either treated or control and refreshed every 48 h, were provided throughout larval development, or for the first week of adult life. Exposure of *C. maculata* larvae to chlorantraniliprole and thiamethoxam applied as seed treatments delayed adult emergence by prolonging the pupal period. When adults were exposed, thiamethoxam reduced the preoviposition

period compared to chlorantraniliprole, whereas the latter treatment cause females to produce fewer clutches during the observation period. Larvae of *C. maculata* did not appear to obtain sufficient hydration from the sunflower stems and their subsequent fecundity and fertility were compromised in comparison to the adult exposure experiment where larvae received supplemental water during development. Exposure of *H. convergens* larvae to thiamethoxam skewed the sex ratio in favor of females; both materials reduced the egg viability of resulting adults and increased the period required for eclosion. Exposure of *H. convergens* adults to chlorantraniliprole reduced egg eclosion times compared to thiamethoxam and exposure to both insecticides reduced pupation times in progeny.

Keywords Biological control Chlorantraniliprole *Coleomegilla maculata* Extrafloral nectar *Hippodamia convergens* Risk assessment Systemic insecticides Thiamethoxam

Introduction

Many beneficial arthropods are omnivorous, consuming both prey and plant material (Coll and Guershon 2002). Natural enemies of insect pests may utilize various plant resources, nibbling tender shoots or consuming pollen and nectar, both floral and extrafloral (Wackers et al. 2007; Lundgren 2009a; Choate and Lundgren 2013). Although floral and extrafloral nectar (EFN) are both rich in sugar, the latter contains sucrose as the dominant sugar, rather than glucose or fructose (Baker and Baker 1979; Rogers 1985). Sugars can be an essential dietary component for coccinellids, improving their survival and reproductive capabilities, and providing metabolic fuel for flight and other behaviors (Lundgren 2009b; Hodek and Evans 2012). Unlike floral nectar which is

available only during flowering, EFN can be available to natural enemies for a much longer period (Pacini et al. 2003; Rose et al. 2006).

On the High Plains of the USA, the EFN secreted by annual sunflowers, *Helianthus annuus* L., (Asteraceae) is an important source of both sugar and hydration for beneficial insects during the hot, dry summers. Although extremely small, the nectaries are highly abundant along petioles and leaf veins and secrete nectar continuously from the time the first true leaves expand until the plants senesce. The production of EFN by plants is associated with attraction of herbivore natural enemies and thus fosters mutualistic protection for the plant (Marazzi et al. 2013). More than 40 species of Coccinellidae are known to utilize EFN in 15 plant families (Pemberton and Vandenberg 1993). A wide variety of insects, beneficial and otherwise, can be directly observed utilizing sunflower EFN a source of hydration during summer months (Charlet and Gavloski 2011) and likely accounts for the great diversity of insects associated with this plant (Royer and Walgenbach 1991).

Coleomegilla maculata DeGeer and *Hippodamia convergens* Guérin-Ménéville (Coleoptera: Coccinellidae) are two of the most abundant lady beetles in the central USA, the latter species being of particular importance for the biological control of cereal aphids in wheat, sorghum and other grains (Rice and Wilde 1988; Nechols and Harvey 1998; Michaud 2013). Both species consume pollen and nectar, both floral and EFN (Pemberton and Vandenberg 1993; Smith and Krischik 1999). These species breed only during periods of high aphid populations, which are usually limited to several weeks in both spring and fall. Aphids and alternative insect prey are rare during summer months, forcing most species into a reproductive diapause (Michaud and Qureshi 2005). Hydration is critical to survival during this period when the beetles must survive on alternative food sources such as pollen and non-aphid prey that have lower water content than aphids, and sunflower EFN can be a key moisture source (Michaud

and Qureshi 2006). Although *C. maculata* can complete development on an exclusive diet of pollen, it has a relatively high water demand when feeding on non-aphid food sources and is sensitive to desiccation stress during development (Michaud and Grant 2005). In contrast, laboratory observations (JP Michaud unpublished) indicate that the daily water consumption of diapausing adult *H. convergens* is only 20-25% that of diapausing *C. maculata*, and the drought tolerance of the former species is likely key to its success in this arid environment.

Recently, both floral nectar and EFN have been recognized as potential routes of exposure to systemic insecticides applied to soil or seeds. Both lethal and sublethal effects have been observed in honey bees (van der Sluijs et al. 2013), predators such as *C. maculata* (Smith and Krischik 1999), *Orius insidiosus* (Say) (Hemiptera: Anthocoridae) (Seagraves and Lundgren 2012; Gontijo et al. 2014a) and *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae) (Rogers et al. 2007; Gontijo et al. 2014b), and parasitoids such as *Anagyrus pseudococci* (Girault) (Hymenoptera: Encyrtidae) (Krischik et al. 2007) and *Lysiphlebus testaceipes* (Cresson) (Hymenoptera: Braconidae) (Moscardini et al. 2014). Seed treatment with systemic insecticides has been widely adopted as a means of early-season pest control in row crops (Hodgson et al. 2012; Nuyttens et al. 2013). However, some studies in soybean, corn and canola have questioned the economic benefit of prophylactic seed treatments, aside from the potential non-target hazards they present (Royer et al. 2005; Wilde et al. 2007; Seagraves and Lundgren 2012). Like most other row crops, the majority of commercial sunflowers are now planted with a systemic insecticide seed treatment, usually thiamethoxam.

The focal insecticides in the present study were chlorantraniliprole and thiamethoxam, examples of two very different insecticide groups, diamides and neonicotinoids, respectively. Both insecticides exhibit systemic activity within

plant vascular tissues which facilitates their use as seed treatments, and their potential to contaminate plant products, including floral and extrafloral nectar (Maienfisch et al. 2001; Lahm et al. 2009; Li et al. 2012). Chlorantraniliprole acts as a ryanodine receptor modulator to block insect muscle contraction; once ingested by an insect, Ca^{++} depletion in muscle cells leads to feeding cessation, lethargy, muscle paralysis and death (Lahm et al. 2007). In contrast, thiamethoxam targets nicotinic acetylcholine receptors in the central nervous system of insects, producing both lethal and sublethal neurological effects (Tomizawa and Casida 2005). The objective of the present study was to assess the sensitivity of *C. maculata* and *H. convergens* to traces of these materials in sunflower EFN and test whether development or reproduction would be impacted when larvae or adults were exposed to sunflower seedlings grown from treated seed.

Materials and methods

Insect colonies

Adults of *C. maculata* and *H. convergens* were collected from fields of sorghum and corn at the Agricultural Research Center in Hays, Kansas, USA (38°51'31.14"N 99°20'10.86"W). Adults of each species were placed in 1-L glass mason jars (ca. 150 per jar) covered with an organdy mesh screen and filled with shredded wax paper as harborage. Water was provided on a cotton wick and approximately 50 mg of frozen *Ephestia kuehniella* (Zeller) (Lepidoptera: Pyralidae) eggs, obtained from a commercial supplier (Beneficial Insectary, Oak Run, CA, USA), were provided daily to each jar. Both species were held in a growth chamber at 24 ± 1 °C, $42 \pm 5\%$ RH, and a photoperiod of

16:8 (L:D). Under these crowded conditions with limited food, the beetles remain in reproductive diapause for many months.

For each experiment, a series of female beetles ($n = 30$) of each species were removed from the jar and isolated, *C. maculata* in plastic Petri dishes (5.5 cm diam) and *H. convergens* in ventilated plexiglass cylinders (5.0 cm diam x 10.0 cm ht) under the same physical conditions as the colony. Females of *C. maculata* were fed with frozen *E. kuehniella* eggs daily with water provided on a small sponge, whereas females of *H. convergens* were fed with an *ad libitum* diet of greenbugs, *Schizaphis graminum* (Rondani) (Hemiptera: Aphididae), because aphids are required to induce oviposition in this species once it has entered diapause (Michaud and Qureshi 2006). The plexiglass cylinders facilitated the provisioning of aphids on excised sorghum seedlings and provided more secure containment of aphids than did the petri dishes. The aphids were obtained from colonies reared on sorghum seedlings in a growth chamber under the same physical conditions as the beetles. Food and water were refreshed daily and eggs, mostly laid on the inner surfaces of the containers, were collected by transferring the beetles to new containers. Upon eclosion, larvae of both species were reared on frozen eggs of *E. kuehniella* in Petri dishes (5.5 cm diam), five per dish, with water provided on a sponge cube, refreshed every 48h, until they emerged as adults. The first laboratory generation was used for larval exposure experiments and the second generation for adult exposure experiments for each species.

Sunflower plants

Triumph Nusun cv. 810CL sunflower seeds were obtained from Triumph Seed Corp. (Dow Agrosciences, Indianapolis, ID) both with and without treatment with Cruiser 5FS[®] (thiamethoxam, 50 mg a.i. 100 kg⁻¹, Syngenta Crop

Protection, Greensboro, NC); untreated Triumph seed served as the experimental control. Sunflower seeds cv. Pioneer 63N82 were obtained from DuPont Crop Protection (E.I. du Pont de Nemours and Co., Wilmington, DE) treated with chlorantraniliprole (1800 mg a.i. 100 kg⁻¹). All seeds were planted 2.0 cm deep in metal trays (8.0 cm x 51.0 cm x 36.0 cm) filled with a mixture of soil, peat moss and perlite (1:1:1) and germinated in a greenhouse at 25 ± 2 °C under natural light supplemented during daylight hours with metal halide lamps (L:D = 12:12). Plants were watered daily, but sparingly, to avoid excessive leaching of insecticide. Sunflower stalks were harvested beginning at the V2 stage (14-15 d-old plants with two true leaves expanded) and every two days thereafter throughout each period of insect exposure, so that insects were exposed to progressively older plant tissues as material was replaced in the experiments. All stem segments were harvested before 9:00 a.m. and constituted the bulk of the upper portion of the main stem. For provisioning to insects, stem segments (ca. 4.0 - 5.0 cm in length) were excised from seedlings and the cut ends dipped in liquid paraffin to seal vascular tissues and maintain turgor, while at the same time preventing the exudation of any resinous materials that might pose a hazard to the insects.

Exposure of larvae

Experiments with both species were conducted under the same physical conditions used for rearing the beetle colonies. Each replicate (n = 8 per treatment) consisted of five first instar larvae held in a Petri dish (5.5 cm diam), their parentage recorded to prevent any subsequent pairing of related beetles. Larvae were fed frozen eggs of *E. kuehniella* ad libitum, refreshed every 48 h. Each Petri dish was supplied with a sunflower stem segment as the only source of hydration, either grown from untreated seed (controls) or from seed treated

with one of the two insecticides. The stem segments were replaced every 48 h until larvae pupated.

Data were recorded daily for all insects throughout the experiment. Larval developmental time was tallied the number of days from the beginning of the experiment until the formation of pupae and pupation time as the number of days from pupal formation to adult emergence. Immature survival was calculated as the percentage of neonate larvae placed in the treatment that successfully emerged as adults. Emergent adults were sexed and, when insects were 7-8 days old, the maximum possible number of pairs were established by confining each female with a male from the same treatment group, checking parentage to prevent the pairing of siblings. After 48 h, males were removed and females were isolated, *C. maculata* in Petri dishes (as above) provisioned with ad libitum frozen eggs of *E. kuehniella* and water on a cube of sponge, *H. convergens* in plexiglass vials (as above) with ad libitum *S. graminum* provided on excised sorghum seedlings. Food and water was refreshed every 48 h.

Eggs were harvested daily by transferring females to new containers and the preoviposition period of each female was calculated as the number of days from adult emergence until first oviposition. A series of 10 clutches were collected from each *H. convergens* female and days required to produce them, recorded. Because oviposition by *C. maculata* was much slower, female fecundity was assessed for a 21 d period post-copula, during which period not all females produced 10 clutches.

Exposure of adults

Pairs of adult beetles (ca. 24 h old) of each species (*C. maculata*, n = 18 per treatment, *H. convergens*, n = 14 per treatment) were established in their respective containers (as above), with parentage checked to prevent the pairing

of siblings. Each container contained a sunflower stem segment corresponding to one of the three treatments and was provisioned with ad libitum frozen eggs of *E. kuehniella*. The sunflower stems were refreshed every 48 h for a total exposure period of 10 days, whereupon males were removed and females isolated for oviposition, those of *C. maculata* receiving eggs of *E. kuehniella* and those of *H. convergens* receiving *S. graminum*. Procedures and data collection were thereafter the same as described above for insects exposed as larvae.

A series of ten neonate larvae hatching from the first clutch of each female were isolated in Petri dishes (5.5 cm diam), one per dish and fed ad libitum frozen eggs of *E. kuehniella* with water provided on a sponge cube, both refreshed every 48h. All insects were observed daily and all developmental data collected until they emerged as adults.

Data analysis

Data were subjected to Kolmogorov-Smirnov and Levine tests ($\alpha = 0.05$) for verification of normality and homoscedasticity, respectively (PROC UNIVARIATE, SAS Institute 2008). For each species and exposure phase (larvae or adults), data that passed these tests were subjected to one-way ANOVA followed and, when means were significant, to a Bonferroni test ($\alpha = 0.05$) to separate means (PROC GLM; SAS Institute 2008). The preoviposition periods of *C. maculata* in the adult exposure test were transformed to $\log(x + 1)$ before being subjected to one-way ANOVA. Untransformed means are presented in all tables. Data that were not normally distributed or failed a Levine test for equality of variances were analyzed using the Kruskal-Wallis test ($\alpha = 0.05$) (PROC NPAR1WAY, SAS Institute 2008). Sex ratio ($\Sigma \text{♀} / \Sigma (\text{♀} + \text{♂})$) was

analyzed using the Chi-square Goodness of Fit test ($\alpha = 0.05$) (PROC FREQ; SAS Institute 2008).

Results

Larvae of *C. maculata* exposed to sunflower stems grown from treated seeds spent about half a day longer in the pupal stage than did controls, but no other treatment effects were significant (Table 1). Survival of pupae was 100% in all three treatments, and the reproductive performance of the resulting females did not differ among treatments, although egg fertility was below normal values in all three. When *C. maculata* were exposed to sunflower stems as adults, there was no mortality in the three week observation period, but preoviposition periods were reduced by the thiamethoxam treatment relative to chlorantraniliprole, although neither was different from controls. However, females in the chlorantraniliprole treatment laid fewer clutches, although overall fecundity and egg viability did not vary significantly among treatments (Table 2). There were no significant treatment effects on any parameter of progeny development.

A higher proportion of emergent adults were female when larvae of *H. convergens* were exposed to stems in the thiamethoxam treatment, but no other developmental parameters differed among treatments (Table 3). Pupal survival was 100% in all three treatments. However, the viability of eggs was significantly reduced for female adults in both the chlorantraniliprole and thiamethoxam treatments and the time required for eclosion of their eggs was slightly increased. Exposure of adults revealed no significant treatment effects on reproductive parameters, but some transgenerational effects were evident in the progeny. Chlorantraniliprole reduced egg eclosion time compared to

thiamethoxam, although controls were not different from either, and both treatments reduced pupation time (Table 4). There was no mortality of adults during the period of reproductive observations.

Discussion

Subtle, but significant, negative effects on development and reproductive biology were observed when *C. maculata* and *H. convergens* fed on EFN presumably contaminated with residues of chlorantraniliprole and thiamethoxam. Systemic insecticides become distributed throughout the plant and may contaminate the pollen, floral and extrafloral nectar (Cloyd and Bethke 2011). EFN can be an important food source for many beneficial organisms, especially coccinellids, because it is rich in sugars that are easily digested (Lundgren 2009b). Consumption of EFN may provide energy and increase fitness, especially when prey is scarce. Lundgren and Seagraves (2011) observed that *C. maculata* adult consuming EFN of *Vicia faba* (Fabaceae) in the absence of prey improved their survival, nutrient reserves and reproductive capacity.

Both seed treatments prolonged *C. maculata* pupation time following larval exposure, possibly because intoxicated larvae had lower mobility and feeding rates, which could have resulted in nutrient limitation and consequent prolongation of the pupal stage or because the materials impaired neural processes controlling pupation. For example, Vargas et al. (2013) showed that *C. maculata* larvae permitted to feed on *E. kuehniella* eggs for only 30 min daily had their total developmental time extended by 10-12 d compared to those permitted ad libitum access to food. Sublethal effects of neonicotinoid insecticides on the foraging behavior and predation rate of beneficial organisms have been previously reported (e.g., Desneux et al. 2007). Smith and Krischik (1999) confined *C. maculata* adults with inflorescences of sunflower plants

treated with imidacloprid via soil and observed significantly reduced motor activity. Imidacloprid reduced the functional response *Serangium japonicum* Chapin (Coleoptera: Coccinellidae) to whitefly *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae) eggs, when applied at a sublethal rate (5 ppm) via egg immersion (He et al. 2012). Thiacloprid applied to tomato at the rate recommended for control of the tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) was shown to cause reductions in the foraging behavior and predation rate of fifth instar nymphs of *Macrolophus pygmaeus* (Hemiptera: Miridae) (Martinou et al. 2014). Thiamethoxam and clothianidin both caused neurotoxic symptoms (e.g., trembling, paralysis, and loss of coordination) in larvae of *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae) exposed for six hours to corn plants grown from treated seeds (Moser and Obrycki 2009). Neonicotinoids also causes negative effects on the motor functions of adult worker honeybees, *Apis mellifera* L. (Hymenoptera: Apidae) (Williamson et al. 2014). Lethargic behavior and feeding inhibition are thus sublethal effects often associated with chlorantraniliprole intoxication. Smagghe et al. (2013) found that workers of *Bombus terrestris* (L.) (Hymenoptera: Apidae) showed lethargic behavior and reduced food consumption following chronic oral exposure to chlorantraniliprole via contaminated pollen.

Chlorantraniliprole has been reported to reduce larval feeding in herbivores such as *Plutella xylostella* L. (Lepidoptera: Plutellidae), *Trichoplusia ni* (Hubner), *Spodoptera exigua* (Hubner) and *Helicoverpa zea* (Boddie) (Lepidoptera: Noctuidae) (Hanning et al. 2009). Oral exposure of neonate *S. exigua* larvae to a sublethal concentration (LC₃₀) of chlorantraniliprole prolonged larval development and increased the appearance of supernumerary instars (Lai and Su 2011). Notwithstanding, the exposure of predatory bugs to surface residues of chlorantraniliprole did not seem to affect foraging behavior in any measurable way. Examples include *Amphiareus*

constrictus (Stal), *Blaptostethus pallescens* Poppius, *Orius tristicolor* (White) (Hemiptera: Anthocoridae) (Pereira et al. 2014), *Podisus nigrispinus* (Dallas) and *Supputius cincticeps* (Stal) (Heteroptera: Pentatomidae) (Castro et al. 2013). Sunflower EFN contaminated with chlorantraniliprole and thiamethoxam caused no lethal effects when consumed by *L. testaceipes* adults, but female foraging behavior was impaired and fewer greenbug nymphs were attacked and parasitized in each bout of foraging (Moscardini et al. 2014). Imidacloprid and chlorantraniliprole also impeded the parasitism of *Nilaparvata lugens* (Stal) (Heteroptera: Delphacidae) by *Anagrus nilaparvatae* (Pang and Wang) (Hymenoptera: Mymaridae) when the wasps consumed honey contaminated with these insecticides (Liu et al. 2010, 2012). Parasitism of aphids by *Aphelinus certus* Yasnosh (Hymenoptera: Aphelinidae) was reduced when the wasps host-fed on aphids that, in turn, fed on imidacloprid- and thiamethoxam-treated soybean plants (Frewin et al. 2014).

Effects of chlorantraniliprole on reproductive biology, similar to those on *C. maculata* in this study, have been reported in other insect groups. For example, Gontijo et al. (2014a) found that female *O. insidiosus* exposed as nymphs to chlorantraniliprole via sunflower EFN suffered extended preoviposition periods. When newly eclosed workers of *B. terrestris* consumed sugar water supplemented with chlorantraniliprole at 40 mg L⁻¹, or pollen sprayed with the insecticide, fewer drones were produced per nest (Smagghe et al. 2013). Negative effects of both insecticides on *H. convergens* biology were delayed and only observed later in life history (reduced egg viability and longer embryonic development when larvae were exposed) or in the next generation (faster pupation time in progeny of exposed adults). In the case of thiamethoxam, increased toxicity is associated with its metabolism into clothianidin, a process which can occur within both plant (Cloyd and Bethke 2011) and insect (Nauen et al. 2003) tissues, and might account for some

delayed impact. Benzidane et al. (2010) reported low toxicity of thiamethoxam to *Periplaneta americana* (L.) (Blattodea: Blattidae) adults was associated with a lack of breakdown into clothianidin within 24 h after its ingestion. In contrast, the primary metabolites of chlorantraniliprole are thought to have low toxicity (FAO, 2008).

Exposure of *H. convergens* larvae to thiamethoxam resulted in a sex ratio skewed toward females when compared to the chlorantraniliprole and control treatments. In contrast, Gontijo et al. (2014b) observed a reduced sex ratio in *C. carnea* when larvae were exposed to sunflower stems grown from seed treated with thiamethoxam as compared to chlorantraniliprole. Thus, patterns of gender-specific larval susceptibility to this material appear to vary among insect groups.

Both larvae and adults of *Coleomegilla maculata* have a high water demand and require supplementary water when feeding on non-aphid foods (Michaud and Grant 2005). The viability of *C. maculata* eggs was abnormally low in the larval exposure experiment, regardless of treatment, where the only moisture available was EFN from the exised stalks, but normal in the adult exposure experiment where larvae were reared with access to water. We infer that the sunflower stalks alone did not provide sufficient hydration for normal larval development in this species, resulting in compromised adult reproduction (reduced fecundity and egg fertility), even though larval survival was relatively good. It is also possible that treatment effects on egg viability, such as those evident in the analogous *H. convergens* experiment, were obscured in these drought-stressed larvae. In *H. convergens*, preoviposition periods averaged two days longer for beetles exposed as adults compared to those exposed as larvae, likely due to the fact the latter received aphid prey two to three days earlier in adult life. Interestingly, both insecticides altered egg eclosion times in *H. convergens* whether the beetles were exposed as larvae or adults, but only the

progeny of *H. convergens* exposed as adults to either insecticide had faster pupation times, with no such effect evident in larvae exposed directly. Thus, the transgenerational effects of these materials do not necessarily mimic the effects of direct exposure.

Transgenerational effects of neonicotinoids have been reported in some coccinellid species. Yu et al. (2014) observed that exposure of *Coccinella septempunctata* L. (Coleoptera: Coccinellidae) larvae to imidacloprid in laboratory microcosms reduced fecundity and egg viability in the next generation. Similar results were reported for topical exposure of *Eriopis connexa* (Gemar) (Coleoptera: Coccinellidae) larvae to acetamiprid that reduced subsequent egg viability (Fogel et al. 2013). Transgenerational effects of systemic insecticides have also been reported for other insect groups, for example thiamethoxam in *Bemisia tabaci* Gennadius (biotype B) and *Trialeurodes vaporariorum* Westwood (Homoptera: Aleyrodidae) (Liang et al. 2012) and chlorantraniliprole in *P. xylostella* (Guo et al. 2013).

Overall, treatment of sunflower seeds with chlorantraniliprole and thiamethoxam caused few negative effects in *C. maculata* and *H. convergens* compared to other beneficial species that have been similarly examined (Gontijo et al. 2014a,b; Moscardini et al. 2014). Because the majority of commercial crops on the High Plains (e.g., sunflower, sorghum, soybeans and corn) are all planted with seed treatments, the beetles colonizing these crops may be regularly exposed to sublethal doses of these insecticides, especially thiamethoxam which has been widely used for more than ten years. Increasing insecticide tolerance in natural enemies, including lady beetles, has been reported as a function of chronic insecticide exposure (Head et al. 1977; Ruberson et al. 2007; Rodrigues et al. 2013a). Rodrigues et al. (2013b) investigated lambda-cyhalothrin susceptibility in 31 populations of lady beetles (Coleoptera: Coccinellidae), focusing on seven species common in cotton fields, and found significant

variation between species and among populations of a given species which they inferred to reflect historical field exposure of the beetles to this insecticide. Thus, the relatively robust responses of both coccinellid species in these experiments may not be representative of other geographic populations that may have different insecticide exposure histories.

In summary, our results suggest that both chlorantraniliprole and thiamethoxam have subtle negative effects on the developmental and reproductive biology of these important predators with the potential for cumulative impacts on population dynamics. Other authors have suggested that the widespread use of systemic insecticides poses a risk to biodiversity and ecosystem services (Biondi et al. 2012; Chagnon et al. 2014; van der Sluijs et al. 2014). The integrated approach to pest management with insecticides is predicated on economic justification and various studies have now failed to identify economic benefits of prophylactic seed treatments in wheat (Royer et al. 2005) corn (Wilde et al. 2007) or soybeans (Seagraves and Lundgren 2012) and others have found them to be ineffective in reducing pests populations (Vernon et al. 2011). Field studies examining a range of nontarget arthropods, conducted over longer time frames, are therefore warranted to determine whether or not these materials are truly compatible with IPM in field crops.

Acknowledgements The authors are grateful to the CAPES Foundation (Brazilian Ministry of Education), the National Council of Scientific and Technological Development (CNPq), and the Minas Gerais State Foundation for Research Aid (FAPEMIG) for scholarship support from CAPES - no. 3362-13-2 (VFM) and CAPES - no. 3363-13-9 (PCG). Voucher specimens are deposited under voucher number 230 in the KSU Museum of Entomological and Prairie Arthropod Research. This is contribution no. 15-260-J of the Kansas State Experiment Station.

Conflict of interest The authors declare that they have no conflicts of interest.

References

- Baker I, Baker HG (1979) Chemical constituents of the nectars of two *Erythrina* species and their hybrid. *Ann Mo Bot Gard* 66:446-450
- Benzidane Y, Touinsi S, Motte E, Jadas-Hecart A, Communal PY, Leduc L, Thany SH (2010) Effect of thiamethoxam on cockroach locomotor activity is associated with its metabolite clothianidin. *Pest Manag Sci* 66:1351-1359
- Biondi A, Desneux N, Siscaro G, Zappala L (2012) Using organic-certified rather than synthetic pesticides may not be safer for biological control agents: selectivity and side effects of 14 pesticides on the predator *Orius laevigatus*. *Chemosphere* 87:803-812
- Castro AA, Correa AS, Legaspi JC, Guedes RN, Serrao JE, Zanuncio JC (2013) Survival and behavior of the insecticide-exposed predators *Podisus nigrispinus* and *Supputius cincticeps* (Heteroptera: Pentatomidae). *Chemosphere* 93:1043-1050
- Chagnon M, Kreutzweiser D, Mitchell EA, Morrissey CA, Noome DA, Van der Sluijs JP (2014) Risks of large-scale use of systemic insecticides to ecosystem functioning and services. *Environ Sci Pollut Res Int*. doi:10.1007/s11356-014-3277-x
- Charlet LD, Gavloski J (2011) Insects of sunflower in the Northern Great Plains of North America. In: Floate KD (ed) *Arthropods of canadian grasslands: inhabitants of a changing landscape*. Biological Survey of Canada, pp 159-178

- Choate BA, Lundgren JG (2013) Why eat extrafloral nectar? Understanding food selection by *Coleomegilla maculata* (Coleoptera: Coccinellidae). *BioControl* 58:359-367
- Cloyd RA, Bethke JA (2011) Impact of neonicotinoid insecticides on natural enemies in greenhouse and interiorscape environments. *Pest Manag Sci* 67:3-9
- Coll M, Guershon M (2002) Omnivory in terrestrial arthropods: mixing plant and prey diets. *Annu Rev Entomol* 47:267-297
- Desneux N, Decourtye A, Delpuech JM (2007) The sublethal effects of pesticides on beneficial arthropods. *Annu Rev Entomol* 52:81-106
- FAO (2008) Food and Agriculture Organization of the United Nations. http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/JMPR/Report08/Chlorantraniliprole.pdf
- Fogel MN, Schneider MI, Desneux N, Gonzalez B, Ronco AE (2013) Impact of the neonicotinoid acetamiprid on immature stages of the predator *Eriopis connexa* (Coleoptera: Coccinellidae). *Ecotoxicology* 22:1063-1071
- Frewin AJ, Schaafsma AW, Hallett RH (2014) Susceptibility of *Aphelinus certus* (Hymenoptera: Aphelinidae) to neonicotinoid seed treatments used for soybean pest management. *J Econ Entomol* 107:1450-1457
- Gontijo PC, Moscardini VF, Michaud JP, Carvalho GA (2014a) Non-target effects of two sunflower seed treatments on *Orius insidiosus* (Hemiptera: Anthocoridae). *Pest Manag Sci*. doi:10.1002/ps.3798
- Gontijo PC, Moscardini VF, Michaud JP, Carvalho GA (2014b) Non-target effects of chlorantraniliprole and thiamethoxam on *Chrysoperla carnea* when employed as sunflower seed treatments. *J Pest Sci* 87:711-719

- Guo L, Desneux N, Sonoda S, Liang P, Han P, Gao X-W (2013) Sublethal and transgenerational effects of chlorantraniliprole on biological traits of the diamondback moth, *Plutella xylostella* L. *Crop Prot* 48:29-34
- Hannig GT, Ziegler M, Marcon PG (2009) Feeding cessation effects of chlorantraniliprole, a new anthranilic diamide insecticide, in comparison with several insecticides in distinct chemical classes and mode-of-action groups. *Pest Manag Sci* 65:969-974
- He Y, Zhao J, Zheng Y, Desneux N, Wu K (2012) Lethal effect of imidacloprid on the coccinellid predator *Serangium japonicum* and sublethal effects on predator voracity and on functional response to the whitefly *Bemisia tabaci*. *Ecotoxicology* 21:1291-1300
- Head R, Neel WW, Sartor CR, Chambers H (1977) Methyl parathion and carbaryl resistance in *Chrysomela scripta* and *Coleomegilla maculate*. *Bull Environ Contam Toxicol* 17:163-164
- Hodek I, Evans EW (2012) Food relationships. In: Hodek I, van Emden HF, Honek A (eds) *Ecology and Behavior of the Ladybird Beetles*. Wiley-Blackwell, Oxford, UK, pp 141-274
- Hodgson EW, Kemis M, Geisinger B (2012) Assessment of Iowa growers for insect pest management practices. *J Extension* 50:RIB6
- Krischik VA, Landmark AL, Heimpel GE (2007) Soil-applied imidacloprid is translocated to nectar and kills nectar-feeding *Anagyrus pseudococci* (Girault) (Hymenoptera: Encyrtidae). *Environ Entomol* 36:1238-1245
- Lahm GP, Cordova D, Barry JD (2009) New and selective ryanodine receptor activators for insect control. *Bioorgan Med Chem* 17:4127-4133
- Lahm GP, Stevenson TM, Selby TP, Freudenberger JH, Cordova D, Flexner L, Bellin CA, Dubas CM, Smith BK, Hughes KA, Hollingshaus JG, Clark CE, Benner EA (2007) Rynaxypyr: a new insecticidal anthranilic

diamide that acts as a potent and selective ryanodine receptor activator.
Bioorg Med Chem Lett 17:6274-6279

- Lai T, Su J (2011) Effects of chlorantraniliprole on development and reproduction of beet armyworm, *Spodoptera exigua* (Hubner). J Pest Sci 84:381-386
- Li X, Degain BA, Harpold VS, Marcon PG, Nichols RL, Fournier AJ, Naranjo SE, Palumbo JC, Ellsworth PC (2012) Baseline susceptibilities of B- and Q-biotype *Bemisia tabaci* to anthranilic diamides in Arizona. Pest Manag Sci 68:83-91
- Liang P, Tian Y-A, Biondi A, Desneux N, Gao X-W (2012) Short-term and transgenerational effects of the neonicotinoid nitenpyram on susceptibility to insecticides in two whitefly species. Ecotoxicology 21:1889-1898
- Liu F, Bao SW, Song Y, Lu HY, Xu JX (2010) Effects of imidacloprid on the orientation behavior and parasitizing capacity of *Anagrus nilaparvatae*, an egg parasitoid of *Nilaparvata lugens*. BioControl 55:473-483
- Liu F, Zhang X, Gui Q-Q, Xu Q-J (2012) Sublethal effects of four insecticides on *Anagrus nilaparvatae* (Hymenoptera: Mymaridae), an important egg parasitoid of the rice planthopper *Nilaparvata lugens* (Homoptera: Delphacidae). Crop Prot 37:13-19
- Lundgren JG (2009a) Relationships of natural enemies and non-prey foods. Springer International, Dordrecht
- Lundgren JG (2009b) Nutritional aspects of non-prey foods in the life histories of predaceous Coccinellidae. Biol Control 5:294-305
- Lundgren JG, Seagraves MP (2011) Physiological benefits of nectar feeding by a predatory beetle. Biol J Linn Soc 104:661-669
- Maienfisch P, Angst M, Brandl F, Fischer W, Hofer D, Kayser H, Kobel W, Rindlisbacher A, Senn R, Steinemann A, Widmer Hr (2001) Chemistry

- and biology of thiamethoxam: a second generation neonicotinoid. *Pest Manag Sci* 57:901-913
- Marazzi B, Bronstein JL, Koptur S (2013) The diversity, ecology and evolution of extrafloral nectaries: current perspectives and future challenges. *Ann Bot* 111:1243-1250
- Martinou AF, Seraphides N, Stavrinides MC (2014) Lethal and behavioral effects of pesticides on the insect predator *Macrolophus pygmaeus*. *Chemosphere* 96:167-173
- Michaud JP (2013) Coccinellids in biological control. In Hodek I, van Emden HF, Honek A (eds) *Ecology and Behaviour of the Ladybird Beetles*. Wiley-Blackwell, Oxford, UK, pp 488-519
- Michaud JP, Grant AK (2005) Suitability of pollen sources for the development and reproduction of *Coleomegilla maculata* (Coleoptera: Coccinellidae) under simulated drought conditions. *Biol Control* 32:363-370
- Michaud JP, Qureshi JA (2005) Induction of reproductive diapause in *Hippodamia convergens* (Coleoptera: Coccinellidae) hinges on prey quality and availability. *Eur J Entomol* 102: 483-487
- Michaud JP, Qureshi JA (2006) Reproductive diapause in *Hippodamia convergens* (Coleoptera: Coccinellidae) and its life history consequences. *Biol Control* 39: 193-200
- Moscardini VF, Gontijo PC, Michaud JP, Carvalho GA (2014) Sublethal effects of chlorantraniliprole and thiamethoxam seed treatments when *Lysiphlebus testaceipes* feed on sunflower extrafloral nectar. *BioControl* 59:503-511
- Moser SE, Obrycki JJ (2009) Non-target effects of neonicotinoid seed treatments; mortality of coccinellid larvae related to zoophytophagy. *Biol Control* 51:487-492

- Nauen R, Ebbinghaus-Kintscher U, Salgado VL, Kausmann M (2003) Thiamethoxam is a neonicotinoid precursor converted to clothianidin in insects and plants. *Pestic Biochem Phys* 76:55-69
- Nechols JR, Harvey TL (1998) Evaluation of a mechanical exclusion method to assess the impact of Russian wheat aphid natural enemies. In Quisenberry SS, Peairs FP (eds) *Response Model for an Introduced Pest – The Russian wheat aphid*. Thomas Say Publications, Lanham, MD, USA, pp 270-279
- Nuyttens D, Devarrewaere W, Verboven P, Foque D (2013) Pesticide-laden dust emission and drift from treated seeds during seed drilling: a review. *Pest Manag Sci* 69:564-575
- Pacini E, Nepi M, Vesprini JL (2003) Nectar biodiversity: a short review. *Plant Syst Evol* 238:7-21
- Pemberton RW, Vandenberg NJ (1993) Extrafloral nectar feeding by ladybird beetles (Coleoptera: Coccinellidae). *P Entomol Soc Wash* 95:139-151
- Pereira RR, Picanço MC, Jr PAS, Moreira SS, Guedes RNC, Corrêa AS (2014) Insecticide toxicity and walking response of three pirate bug predators of the tomato leaf miner *Tuta absoluta*. *Agr Forest Entomol*. doi:10.1111/afe.12059
- Rice ME, Wilde GE (1988) Experimental evaluation of predators and parasitoids in suppressing greenbugs (Homoptera: Aphididae) in sorghum and wheat. *Environ Entomol* 17:836-841
- Rodrigues ARS, Ruberson JR, Torres JB, Siqueira HÁA, Scott JG (2013a) Pyrethroid resistance and its inheritance in a field population of *Hippodamia convergens* (Guérin-Ménéville) (Coleoptera: Coccinellidae). *Pestic Biochem Phys* 105:135-143
- Rodrigues ARS, Spindola AF, Torres JB, Siqueira HA, Colares F (2013b) Response of different populations of seven lady beetle species to

- lambda-cyhalothrin with record of resistance. *Ecotox Environ Safe* 96:53-60
- Rogers CE (1985) Extrafloral nectar: entomological implications. *Bull Entomol Soc Am* 31:15-20
- Rogers MA, Krischik VA and Martin LA (2007) Effect of soil application of imidacloprid on survival of adult green lacewing, *Chrysoperla carnea* (Neuroptera: Chrysopidae), used for biological control in greenhouse. *Biol Control* 42:172-177
- Rose USR, Lewis J, Tumlinson JH (2006) Extrafloral nectar from cotton (*Gossypium hirsutum*) as a food source for parasitic wasps. *Funct Ecol* 20:67-74
- Royer TA, Giles KL, Nyamanzi T, Hunger RM, Krenzer EG, Elliott NC, Kindler SD, Payton M (2005) Economic evaluation of the effects of planting date and application rate of imidacloprid for management of cereal aphids and barley yellow dwarf in winter wheat. *J Econ Entomol* 98:95-102
- Royer TA, Walgenbach DD (1991) Predacious arthropods of cultivated sunflower in eastern South Dakota. *J Kansas Entomol Soc* 64:112-116
- Ruberson JR, Roberts P, Michaud JP (2007) Pyrethroid resistance in Georgia populations of the predator *Hippodamia convergens* (Coleoptera: Coccinellidae). *Proc Beltwide Cotton Conf* 1:361-365
- SAS Institute (2008) SAS for Windows Version 9.0. SAS Institute. Cary, North Carolina
- Seagraves MP, Lundgren JG (2012) Effects of neonicotinoid seed treatments on soybean aphid and its natural enemies. *J Pest Sci* 85:125-132
- Smagghe G, Deknopper J, Meeus I, Mommaerts V (2013) Dietary chlorantraniliprole suppresses reproduction in worker bumblebees. *Pest Manag Sci* 69:787-791

- Smith SF, Krischik VA (1999) Effects of systemic imidacloprid on *Coleomegilla maculata* (Coleoptera: Coccinellidae). *Environ Entomol* 28:1189-1195
- Tomizawa M, Casida JE (2005) Neonicotinoid insecticide toxicology: mechanisms of selective action. *Annu Rev Pharmacol* 45:247-268
- van der Sluijs JP, Amaral-Rogers V, Belzunces LP, Lexmond MFIJBv, Bonmatin J-M, Chagnon M, Downs CA, Furlan L, Gibbons DW, Giorio C, Girolami V, Goulson D, Kreuzweiser DP, Krupke C, Liess M, Long E, McField M, Mineau P, Mitchell EAD, Morrissey CA, Noome DA, Pisa L, Settele J, Simon-Delso N, Stark JD, Tapparo A, Dyck HV, Praagh Jv, Whitehorn PR, Wiemers M (2014) Conclusions of the worldwide integrated assessment on the risks of neonicotinoids and fipronil to biodiversity and ecosystem functioning. *Environ Sci Pollut Res*. doi:10.1007/s11356-014-3229-5
- van der Sluijs JP, Simon-Delso N, Goulson D, Maxim L, Bonmatin J-M, Belzunces LP (2013) Neonicotinoids, bee disorders and the sustainability of pollinator services. *Curr Opin Environ Sustainability* 5:293-305
- Vargas G, Michaud JP, Nechols JR (2013) Trajectories of reproductive effort in *Coleomegilla maculata* and *Hippodamia convergens* (Coleoptera: Coccinellidae) respond to variation in both income and capital. *Environ Entomol* 42:341-353
- Vernon RS, Herk WG, Clodius M, Harding C (2011) Crop protection and mortality of *Agriotes obscurus* wireworms with blended insecticidal wheat seed treatments. *J Pest Sci* 86:137-150
- Wackers FL, Romeis J, van Rijn P (2007) Nectar and pollen feeding by insect herbivores and implications for multitrophic interactions. *Annu Rev Entomol* 52:301-323

- Wilde G, Roozeboom K, Ahmad A, Claassen M, Gordon B, Heer W, Maddux L, Martin V, Evans P, Kofoid K, Long J, Schlegel A, Witt M (2007) Seed treatment effects on early season pests of corn and corn growth and yield in the absence of agricultural pests. *J Agr Urban Entomol* 24:177-193
- Williamson SM, Willis SJ, Wright GA (2014) Exposure to neonicotinoids influences the motor function of adult worker honeybees. *Ecotoxicology* 23:1409-1418
- Yu C, Lin R, Fu M, Zhou Y, Zong F, Jiang H, Lv N, Piao X, Zhang J, Liu Y, Brock TCM (2014) Impact of imidacloprid on life-cycle development of *Coccinella septempunctata* in laboratory microcosms. *Ecotox Environ Safe* 110:168-173

Table 1 Mean (\pm SE) developmental and reproductive parameters of *Coleomegilla maculata* exposed as larvae to sunflower stems grown from treated seeds.

Parameter	Seed treatment			<i>F</i> , <i>H</i> or χ^2	df	<i>P</i>
	Untreated	Chlorantraniliprole	Thiamethoxam			
Larval survival (%)	72.5 \pm 8.4	70.0 \pm 7.6	87.5 \pm 6.5	3.06*	2	0.216
Larval development time (d)	11.7 \pm 0.2	11.7 \pm 0.3	11.9 \pm 0.2	0.13	2,21	0.881
Pupation time (d)	3.5 \pm 0.1 b	3.9 \pm 0.1 a	3.9 \pm 0.1 a	8.89	2,21	0.002
Sex ratio ¹ (n)	0.55 \pm 0.09 (29)	0.50 \pm 0.09 (28)	0.40 \pm 0.08 (35)	1.54**	2	0.463
No. pairs mated	9	10	12			
Preoviposition period (d)	14.8 \pm 2.1	18.1 \pm 2.1	16.0 \pm 1.6	0.72	2,28	0.494
Fecundity (eggs female ⁻¹)	102.0 \pm 29.1	94.6 \pm 31.6	122.1 \pm 24.9	0.78*	2	0.677
No. clutches laid in 21 d	8.3 \pm 2.1	7.3 \pm 1.9	9.5 \pm 1.1	0.44	2,28	0.650
Egg viability (% hatching)	50.7 \pm 7.1	36.9 \pm 6.3	37.5 \pm 6.7	1.23	2,28	0.307
Eclosion time (d)	3.0 \pm 0.0	3.0 \pm 0.0	3.1 \pm 0.1	2.55	2,25	0.098

Analysis by one-way ANOVA, Kruskal-Wallis (*) or Chi-square (**).

Means followed by different letters were significantly different within rows (Bonferroni test, $\alpha = 0.05$).

¹proportion female.

Table 2 Mean (\pm SE) reproductive parameters of *Coleomegilla maculata* adults exposed to sunflower stems grown from treated seed and the developmental parameters of their offspring.

Parameter	Seed treatment			<i>F</i> , <i>H</i> or χ^2	df	<i>P</i>
	Untreated	Chlorantraniliprole	Thiamethoxam			
No. pairs established	18	18	18			
Preoviposition period (d)	14.3 \pm 1.1 ab	16.9 \pm 1.6 a	12.4 \pm 0.6 b	3.62	2,51	0.034
Fecundity (eggs female ⁻¹)	188.1 \pm 21.8	152.8 \pm 29.6	210.8 \pm 26.4	1.25	2,51	0.294
No. clutches laid in 21 d	14.1 \pm 1.1 a	10.4 \pm 1.6 b	15.1 \pm 1.0 a	3.79	2,51	0.029
Egg viability (% hatching)	92.1 \pm 2.3	88.9 \pm 1.8	83.9 \pm 5.5	1.30	2,51	0.283
Eclosion time (d)	3.0 \pm 0.0	3.1 \pm 0.0	3.0 \pm 0.1	0.79*	2	0.673
Larval development time (d)	11.7 \pm 0.1	12.0 \pm 0.1	11.9 \pm 0.1	2.28	2,51	0.113
Pupation time (d)	3.1 \pm 0.0	3.1 \pm 0.1	3.1 \pm 0.1	0.19	2,51	0.828
Immature survival (%)	96.1 \pm 1.2	96.7 \pm 1.9	94.4 \pm 1.8	1.97*	2	0.373
Sex ratio ¹ (n)	0.57 \pm 0.04 (173)	0.53 \pm 0.04 (171)	0.55 \pm 0.04 (170)	0.74**	2	0.692

Analysis by one-way ANOVA, Kruskal-Wallis (*) or Chi-square (**).

Means followed by different letters were significantly different within rows (Bonferroni test, $\alpha = 0.05$).

¹proportion female.

Table 3 Mean (\pm SE) developmental and reproductive parameters of *Hippodamia convergens* exposed as larvae to sunflower stems grown from treated seeds.

Parameter	Seed treatment			<i>F</i> or χ^2	df	<i>P</i>
	Untreated	Chlorantraniliprole	Thiamethoxam			
Larval survival (%)	67.5 \pm 6.5	67.5 \pm 6.4	57.5 \pm 10.9	0.49	2,21	0.620
Larval development time (d)	11.7 \pm 0.2	12.0 \pm 0.2	11.9 \pm 0.3	0.55	2,20	0.587
Pupation time (d)	4.8 \pm 0.1	4.6 \pm 0.1	4.8 \pm 0.1	0.86	2,20	0.448
Sex ratio ¹ (n)	0.33 \pm 0.09 (27) b	0.31 \pm 0.09 (26) b	0.61 \pm 0.10 (23) a	6.54*	2	0.043
No. pairs mated	9	7	8			
Preoviposition period (d)	9.6 \pm 1.1	10.3 \pm 1.3	8.9 \pm 1.5	0.27	2,21	0.767
Fecundity (eggs female ⁻¹)	251.3 \pm 19.4	262.1 \pm 23.0	254.0 \pm 27.7	0.05	2,21	0.947
No. days for 10 clutches	10.6 \pm 0.2	11.3 \pm 0.5	11.5 \pm 0.9	0.76	2,21	0.479
Egg viability (% hatching)	93.2 \pm 1.9 a	79.9 \pm 4.1 b	84.1 \pm 3.4 b	4.87	2,21	0.018
Eclosion time (d)	3.1 \pm 0.0 b	3.3 \pm 0.0 a	3.2 \pm 0.0 a	4.56	2,21	0.023

Analysis by one-way ANOVA or Chi-square (*).

Means followed by different letters were significantly different within rows (Bonferroni test or Chi-square, $\alpha = 0.05$).

¹proportion female.

Table 4 Mean (\pm SE) reproductive parameters of *Hippodamia convergens* adults exposed to sunflower stems grown from treated seed and the developmental parameters of their offspring.

Parameter	Seed treatment			<i>F</i> , <i>H</i> or χ^2	df	<i>P</i>
	Untreated	Chlorantraniliprole	Thiamethoxam			
No. pairs mated	14	14	14			
Preoviposition period (d)	12.1 \pm 0.7	12.4 \pm 0.6	13.2 \pm 0.3	1.17	2,39	0.332
Fecundity (eggs female ⁻¹)	247.6 \pm 16.7	287.7 \pm 18.8	264.1 \pm 18.4	1.27	2,39	0.294
No. days for 10 clutches	13.6 \pm 0.6	12.1 \pm 0.6	12.5 \pm 0.4	2.38	2,39	0.106
Egg viability (% hatching)	86.7 \pm 2.8	82.8 \pm 2.8	79.3 \pm 5.7	1.02*	2	0.602
Eclosion time (d)	3.2 \pm 0.0 ab	3.1 \pm 0.0 b	3.3 \pm 0.0 a	3.27	2,39	0.049
Larval development time (d)	11.5 \pm 0.1	11.7 \pm 0.1	11.7 \pm 0.1	1.22	2,38	0.308
Pupation time (d)	4.9 \pm 0.1 a	4.7 \pm 0.1 b	4.7 \pm 0.1 b	4.55	2,38	0.017
Immature survival (%)	86.4 \pm 2.9	85.0 \pm 4.5	83.8 \pm 2.9	0.77*	2	0.682
Sex ratio ¹ (n)	0.53 \pm 0.05 (121)	0.45 \pm 0.05 (119)	0.46 \pm 0.05 (109)	2.09**	2	0.353

Analysis by one-way ANOVA, Kruskal-Wallis (*) or Chi-square (**).

Means followed by different letters were significantly different within rows (Bonferroni test, $\alpha = 0.05$).

¹proportion female.